## DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

T. B. C.: DS - 2015

Serial No.

1783

**Test Booklet Series** 



## TEST BOOKLET

**DENTAL SURGEON** 

Time Allowed : 3 Hours

Maximum Marks : 200

## : INSTRUCTIONS TO CANDIDATES :

- 1. IMMEDIATELY AFTER THE COMMENCEMENT OF THE EXAMINATION, YOU SHOULD CHECK THAT THIS TEST BOOKLET **DOES NOT** HAVE ANY UNPRINTED OR TORN OR MISSING PAGES OR ITEMS ETC. IF SO, GET IT REPLACED BY A COMPLETE TEST BOOKLET OF THE SAME SERIES ISSUED TO YOU.
- 2. ENCODE CLEARLY THE TEST BOOKLET SERIES **A**, **B**, **C** OR **D**, AS THE CASE MAY BE, IN THE APPROPRIATE PLACE IN THE ANSWER SHEET USING BALL POINT PEN (BLUE OR BLACK).
- 3. You have to enter your Roll No. on the Test Booklet in the Box provided alongside. DO NOT write anything else on the Test Booklet.
- 4. This Test Booklet contains 200 items (questions). Each item (question) comprises four responses (answers). You have to select the correct response (answer) which you want to mark (darken) on the Answer Sheet. In case, you feel that there is more than one correct response (answer), you should mark (darken) the response (answer) which you consider the best. In any case, choose ONLY ONE response (answer) for each item (question).
- You have to mark (darken) all your responses (answers) ONLY on the separate Answer Sheet provided, by using BALL POINT PEN (BLUE OR BLACK). See instructions in the Answer Sheet.
- 6. All items (questions) carry equal marks. All items (questions) are compulsory. Your total marks will depend only on the number of correct responses (answers) marked by you in the Answer Sheet. There will be no negative marking for wrong answer.
- 7. Before you proceed to mark (darken) in the Answer Sheet the responses to various items (questions) in the Test Booklet, you have to fill in some particulars in the Answer Sheet as per the instructions sent to you with your **Admission Certificate**.
- 8. After you have completed filling in all your responses (answers) on the Answer Sheet and after conclusion of the examination, you should hand over to the Invigilator the *Answer Sheet* issued to you. You are allowed to take with you the candidate's copy/second page of the Answer Sheet along with the *Test Booklet* after completion of the examination for your reference.

DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

1.		weakest part of mandible where ure occurs is :	6.	-	es, stones moans and groans
	(A) (B) (C) (D)	Neck of condyle Angle of mandible Midline Canine fossa		(A) (B) (C) (D)	Hyperparathyroidism Hypoparathyroidism Hypoparathyroidism Hypothyroidism
2.	Cleft (A) (B) (C) (D)	palate repair is ideally done at : Birth 1-6 months 7-18 months 24-36 months	7.		suture that maintains strength for est time is: Polyglycolic Acid (Dexon) Polyglactin (Vicryl) Polydioxone (PDS) Polyglyconate (Chromic catgut)
3.	fracti (A) (B) (C) (D)	most commonly involved bone in ure of the face is :  Nasal  Maxilla  Mandible  Zygoma	8.		min K is required by liver for the luction of active : Fibrinogen Platelet Globulins Prothrombin
4.		inoma of the lower alveolar coccurs around : Incisor Canine Molar Pre molar	9.	arte	itation of baroreceptors in ries reflexly causes : Peripheral vasoconstriction Arterial pressure to decrease Increase cardiac output Vasovagal syncope
5.		vary stones occurs most monly in : Parotid Sublingual Submandibular Submaxillary	10.	Card of: (A) (B) (C) (D)	Stroke volume × venous return Stroke volume × heart rate Venous return × heart rate Stroke volume × cardiac index
IB-	(C) 1C/34	· .	<b>)</b> .	(D)	Contd.

11.	Pre-	optic area of hypothalamus is	16.	Pain	sensation travels through the:
	cond	erned with:		(A)	Posterior column
	(A)	Milk ejection		(B)	Lateral spinothalamic tract
	(B)	Uterine contractility		(C)	Corticospinal tract
	(C)	Temperature regulation		(D)	Vestibulospinal tract
*	(D)	Memory	17.	ADH	I increases the permeability of
12.	REM	A sleep is associated with :		dista	al tubule to :
	(A)	Normal heart rate		(A)	Proteins
	(B)	Increased muscle tone		(B)	Amino acids
	(C)	Decreased respiratory rate		(C) <sub>.</sub>	Glucose
	(D)	Extreme inhibition of peripheral		(D)	Water
		muscles	18.	Hae	moprotein which binds to oxygen
13.	Sym	pathetic stimulation :			t strongly is :
	(A)	Increases gastric motility		(A)	Adult haemoglobin
	(B)	Causes Pupillary dilation		(B)	Foetal haemoglobin
	(C)	Decreases cardiac activity		(C)	Haemosiderin
	(D)	Vasodilation		(D)	Myoglobin
14.		ch of the following best describes Broca area of speech?	19.		iciency of which one of the wing does not lead to anemia?
	(A)	Sensory speech area		(A)	Serum Iron
	(B)	Controls emotions		(B)	Serum Copper
	(C)	Located in temporal bone		(C)	Vitamin C
	(D)	Motor speech area		(D)	Serum Zinc
15.		entional tremor occurs in the ease of:	20.		st of the enzymes secreted in the nan gastrointestinal tract are :
3]:	(A)	Cerebrum		(A)	Oxidases
	(B)	Basal ganglia		(B)	Reductases
	(C)	Cerebellum		(C)	Lyases
	(D)	Spinal cord		(D)	Hydrolases
IR -	· 1C/3		(3)	` '	(Turn over)
		· •			•

- 21. The primary function of the salivary amylase is:
  - (A) Hydrolysis of dietary starch
  - (B) Hydrolysis of bacterial cell wall
  - (C) Hydrolysis of glycogen
  - (D) Hydrolysis of pectin
- 22. Final common oxidative pathway which integrates oxidative products of fats, proteins and carbohydrates is:
  - (A) Gluconeogenesis
  - (B) Glycolysis
  - (C) Citric acid cycle
  - (D) Anaerobic glycolysis
- 23. Largest amount of energy is produced by which of the following?
  - (A) Oxidation of amino acids
  - (B) Beta oxidation of saturated fatty acids
  - (C) Beta oxidation of unsaturated fatty acid
  - (D) Anaerobic glycolysis
- 24. The substance that contributes maximally to the osmolality inside the cell is:
  - (A) Protein
  - (B) Phosphate
  - (C) Sodium
  - (D) Urea

- 25. The unique feature of Mitochondria is its:
  - (A) Actin
  - (B) Myosin
  - (C) Prothrombin
  - (D) DNA
- 26. Calcifying epithelial odontogenic tumour commonly occurs in :
  - (A) Mandibular Posterior area
  - (B) Maxillary Posterior area
  - (C) Mandibular Anterior area
  - (D) Maxillary Anterior area
- 27. The most common histological variant of Ameloblastoma is:
  - (A) Follicular
  - (B) Plexiform
  - (C) Acanthomatous
  - (D) Granular cell
- 28. The histological variant of ameloblastoma with highest recurrenence rate is:
  - (A) Follicular
  - (B) Acanthomatous
  - (C) Granular cell
  - (D) Basal cell
- 29. Smoking is associated with the genesis of which of the salivary gland tumor?
  - (A) Pleomorphic adenoma

IB - 1C/34

(4)

- (B) Warthin's tumor
  (C) Necrotizing sialometaplasia
  (D) Papillary cystadenoma
  30. A tumor which arises in salivary gland tissue and entrapped in parotid lymph nodes during embryogenesis is:
  (A) Pleomorphic adenoma
  (B) Warthin's tumor
  (C) Oncocytoma
  (D) Mucinous adenocarcinoma
- 31. In which of the following condition the normal flora becomes pathogenic?
  - (A) Diabetes
  - (B) Leukemia
  - (C) AIDS
  - (D) All of the above
- 32. Following are the beneficial effects of the resident flora except:
  - (A) Synthesis of Vitamin K
  - (B) Production of Pencillinase
  - (C) Suppression of colonization by pathogen
  - (D) Bacteriocin production
- 33. Which of the following can be a congenital infection?
  - (A) AIDS
  - (B) Toxoplasmosis
  - (C) CMV
  - (D) All of the above

- 34. Following bacteria produces enterotoxin except:
  - (A) Staph aureus
  - (B) B. anthracis
  - (C) B. cereus
  - (D) V cholerae
- 35. Which of the following disease is not acquired by ingestion?
  - (A) Salmonellosis
  - (B) Shigellosis
  - (C) Diptheria
  - (D) Cholerae
- 36. Following bacteria are invasive except:
  - (A) Shigella
  - (B) S. Typhi
  - (C) Vibiro Cholerae
  - (D) N. Meningitides
- 37. Which of the following is vector borne disease?
  - (A) Typhus
  - (B) Dengue
  - (C) Yellow fever
  - (D) All of the above
- 38. Plasmid mediated resistance was first detected in :
  - (A) Salmonella
  - (B) Shigella
  - (C) Proteus
  - (D) E. coli

(Turn over)

39.	Following are zoonotic of except:	lisease 44	. Ea	mes technique provides for :
	(A) Rabies	•	(A)	Low mercury-alloy ratio
	(B) Toxoplasmosis		(B)	High mercury-alloy ratio
	(C) Hydatid		(C)	Easy waste management
	(D) Tetanus		(D)	Good polishing
40.		<del>-</del> .	esti	rogenic material?
	•		(A)	Bisphenol – A
	(A) Transcription (B) Translation		(B)	Poly acrylic acid
	` ,		(C)	Nickel
	(C) Conjugation (D) Transduction		(D)	DFG
41.	(D) Transduction  Which is the cement with the solubility?	· 46. highest		nultaneous application of acid to mel and dentine is known as :
	(A) Ca(OH) <sub>2</sub>		(A)	Total etch technique
	(B) Silicate		(B)	Dry bonding
	(C) $Zn(PO_{d})$		(C)	Wet bonding
	(D) GIC		(D)	Hybridization
42.	Advantage of LED ligth is:	47.	Sme	ear layer consists of :
	(A) Small and portable		(A)	Enamel debris
	(B) Low power consumption	า	(B)	Microorganism
	(C) Minimal deterioration of	of light	(C)	Dentin chips
	output (D) All of the above		(D)	All of the above
43.	(D) All of the above  The angle between rake factorized radial line is:	e and 48.	Cem pulp	nent which is kindest to dental
			(A)	GIC
	(A) Rake angle		(B)	Polycarboxylate
	(B) Clearance angle		` '	•
	(C) Blade angle		(C)	Silicate
	(D) Radial clearance		(D)	Resin cement
IB –	1C/34	(6)		Contd.

49.	Cori	rugated gold is a type of :			(C)	Hepatitis C
	(A)	Casting alloy			(D)	Hepatitis D
	(B)	DFG		54.	Bloc	od Coagulation Factor which is
	(C)	Technic alloy				required in the intrinsic pathway
	(D)	None of the above			for c	coagulation is :
50.	Elas	stic modulus of a material	1		(A)	Factor 9
	indic	cates its :			(B)	Factor 11
	(A)	Rigidity			(C)	Factor 7
	(B)	Flexibility			(D)	Factor 13
	(C)	Proportional limit		55.	All t	he following are suggestive of
	(D)	Viscosity			iron	deficiency anemia except :
51.	Swa	n-neck deformity of the fingers			(A)	Koilonychia
	is se	en in :			(B)	Low reticulocyte response
	(A)	Osteoarthritis			(C)	Decreased serum ferritin
	(B)	Rheumatoid arthritis			(D)	Decreased Total Iron-binding
	(C)	Osteomalacia				Capacity (TIBC)
	(D)	Osteoporosis		56.	Alla	re side effects of corticosteroids
52.	Arth	us type allergic reaction is a			exce	ept:
	featu	re of :			(A)	Myopathy
	(A)	Type I reactions			(B)	Diarrhoea
	(B)	Type II reactions			(C)	Peptic ulcer disease
	(C)	Type III reactions			(D)	Osteoporosis
	(D)	Type IV reactions		57.	Vitar	min B12 deficiency may cause
53.	Prog	ression to chronic liver disease			all of	the following except:
		een most commonly with			(A)	Optic neuritis
	infed	tion by:			<b>(B)</b>	Myelopathy
	(A)	Hepatitis A			(C)	Myopathy
	(B)	Hepatitis B			(D)	Neuropathy
B-	1C/34	ļ.	(7)		•	(Turn over)

dependent coagulation factors except:  (A) Factor X  (B) Factor IX  (C) Factor III  (D) Factor VII  (A) Roof of pasopharynx  (B) Osteopetrosis  (C) Osteoporosis  (D) Osteonecrosis				•
(C) Radius (D) Vertebra (A) ACTH stimulation test (B) Plasma Cortisol (C) Plasma FSH (D) Plasma Aldosterone (C) Plasma FSH (D) Plasma Aldosterone (E) Plasma Cortisol (C) Plasma FSH (D) Plasma Aldosterone (E) Plasma FSH (D) Plasma FSH (D) Plasma Aldosterone (E) Plasma FSH (D) Plasma Serological marker is suggestive of highly infectious stage of acute Hepatitis B? (A) HBsAg (B) Anti HBsAg (C) HBeAg (C) HBeAg (D) Anti HBeAg (D) Anti HBeAg (E) Normal Serum Calcium, Phosphate and ALP are seen in: (A) Osteomalacia (B) Osteopetrosis (C) Osteoporosis (D) Osteonecrosis (E) Osteoporosis (D) Osteonecrosis (E) Nasopharyngeal carcinoma mostly arises from the: (A) Roof of nasopharynx (B) Posterior wall	58.	common site for a fracture associated with osteoporosis?		(C) Cushing Syndrome
59. All are the causes of bilateral parotid enlargement except:  (A) Diabetes mellitus (B) Systemic lupus erythromatosus (C) HIV suggestive of highly infectious stage of acute Hepatitis B?  60. Which of the following is not a clinical feature of cerebellar disease? (A) Nystagmus (B) Hyperreflexia (C) Hypotonia (D) Tremors  61. All the following are Vitamin K-dependent coagulation factors except: (A) Factor X (B) Factor III (D) Factor VII  62. Excessive weight gain, menorrhagia, cold intolerance and weakness are suggestive of highly infectious stage of acute Hepatitis B?  (A) HBsAg (B) Anti HBsAg (C) HBeAg (D) Anti HBeAg (D) Anti HBeAg (E) Normal Serum Calcium, Phosphate and ALP are seen in: (A) Osteomalacia (B) Osteopetrosis (C) Osteoporosis (D) Osteonecrosis (C) Ansopharyngeal carcinoma mostly arises from the: (A) Roof of nasopharynx (B) Posterior wall (C) Anterior wall		(C) Radius	63.	of Cushing syndrome is :  (A) ACTH stimulation test
(C) HIV suggestive of highly infectious stage of acute Hepatitis B?  60. Which of the following is not a clinical feature of cerebellar disease?  (A) Nystagmus (B) Hyperreflexia (C) Hypotonia (D) Tremors  61. All the following are Vitamin K-dependent coagulation factors except: (A) Factor X (B) Factor IX (C) Factor III (D) Factor VII  62. Excessive weight gain, menorrhagia, cold intolerance and weakness are suggestive of:  (A) Which of the following is not a clinical of acute Hepatitis B?  (A) HBsAg (B) Anti HBsAg (C) HBeAg (D) Anti HBeAg (E) Normal Serum Calcium, Phosphate and ALP are seen in: (A) Osteomalacia (B) Osteopetrosis (C) Osteoporosis (D) Osteonecrosis (D) Osteonecrosis (E) Nasopharyngeal carcinoma mostly arises from the: (A) Roof of nasopharynx (B) Posterior wall (C) Anterior wall	59.	enlargement except :		(C) Plasma FSH
feature of cerebellar disease?  (A) Nystagmus (B) Hyperreflexia (C) Hypotonia (D) Tremors  61. All the following are Vitamin K-dependent coagulation factors except: (A) Factor X (B) Factor IX (C) Factor III (D) Factor VII  62. Excessive weight gain, menorrhagia, cold intolerance and weakness are suggestive of:  (B) Anti HBsAg (C) HBeAg (D) Anti HBeAg (E) HBeAg (D) Anti HBsAg (E) HBeAg (E) HBeA		(C) HIV	64.	suggestive of highly infectious stage
(D) Tremors  65. Normal Serum Calcium, Phosphate and ALP are seen in:  (A) Osteomalacia (B) Osteopetrosis (C) Osteoporosis (D) Osteonecrosis (D) Osteonecrosis (E) Osteoporosis (D) Osteoporosis (E) Osteoporosis (	60.	feature of cerebellar disease?  (A) Nystagmus		(B) Anti HBsAg (C) HBeAg
dependent coagulation factors except:  (A) Factor X (B) Factor IX (C) Factor III (D) Factor VII  62. Excessive weight gain, menorrhagia, cold intolerance and weakness are suggestive of:  (B) Osteopetrosis (C) Osteoporosis (D) Osteonecrosis  68. Nasopharyngeal carcinoma mostly arises from the: (A) Roof of nasopharynx (B) Posterior wall (C) Anterior wall			65.	•
(D) Factor VII arises from the :  (A) Roof of nasopharynx  cold intolerance and weakness are suggestive of :  (C) Anterior wall	61.	dependent coagulation factors except:  (A) Factor X		(B) Osteopetrosis (C) Osteoporosis
cold intolerance and weakness are suggestive of:  (B) Posterior wall  (C) Anterior wall			66.	arises from the :
/= /	62.	cold intolerance and weakness are suggestive of:		(B) Posterior wall (C) Anterior wall

67.	_	nest incidence of naso- yngeal carcinoma is seen in :	72.	Clef age	t lip is ideally repaired at what	
	(A)	Indian	•	(A)	Soon after birth	
	(B)	Chinese		(B)	6-9 weeks	
	(C)	Americans		(C)	10-24 weeks	
	(D)	Europeans		(D)	25-36 weeks	
68.	Whi	ch region is most commonly	73.	Den	tigerous cyst develops around :	
-	affe	cted in actinomycosis?		(A)	Root	
	(A)	A) Thorax	•	(B)	Crown	
	(B)	Liver		(C)	Gum	
	(C)	Facio-cervical		(D)	Gingival Margin	
	(D)	Liver				
69.	A punched out edge is characteristic of which ulcer?		74.		In tongue cancers, the least affected	
				site	•	
	(A)	Tuberculosis		(A)	Ventral margin	
	(B)	Malignant		(B)	Dorsal margin	
	(C)	Rodent		(C)	Тір	
	(D)	Syphilis		(D)	Lateral margin	
70.	Cystic hygroma is a :		75.	Ame	Ameloblastoma most commonly	
, 0.	(A)	Dermoid cyst		deve	elops around :	
	(B)	Cavernous haemangioma		(A)	Incisors	
	(C)	Lymphangiectasia		(B)	Premolars	
	(D)	Sebaceous cyst		(C)	Canine	
				(D)	Third molar	
71.		most common indication for the	76	Man	y substances are removed from	
	remo	oval of sublingual salivary gland			cell to outside by :	
	(A)	Ranula		(A)	Phagocytosis	
	(B)	Lymphangioma		(B)	Pinocytosis	
	(C)	Neoplasm		(C)	Exocytosis	
	(D)	Stone		(D)	Chemotaxis	
			7.03	(2)		
IB -	1C/3	4	(9)		(Turn over)	

77.	Ner	ve fibre with slowest conduction	82.		ich of the following is a branch of killary nerve ?	
	(A) (B)	A alpha A gamma		(A) (B)	Supratrochlear  Zygomaticotemporal	
	(C)	Beta fiber		(C)	Infratrochlear	
	(D)	C fibres		(D)	External nasal	
78.	The superior meatus of the nose contains the opening of:		83.		e taste sensation from the terior one-third of the tongue is	
	(A)	Anterior ethmoidal air cells		carr	ied by :	
	(B)	Middle ethmoidal air cells		(A)	Chorda tympani	
	(C)	Posterior ethmoidal air cells		(B)	Lingual nerve	
	(D)	Maxillary air sinus		(C)	Hypoglossal nerve	
79.	The	posterior auricular nerve is a		(D)	Glossopharyngeal nerve	
	branch of:		84.	Which of the following muscle assists		
	(A)	Chorda tympani		in th	e elevation of mandible ?	
	(B)	Facial		(A)	Medial pterygoid	
	(C)	Auriculo temporal		(B)	Lateral pterygoid	
	(D)	Accessory nerve		(C)	Digastric	
80.	The	lateral wall of the infratemporal		(D)	Genohyoid	
	fossa is formed by:		85.	The	The sensory root of the facial nerve	
	(A)	Lateral pterygoid plate		is known as :		
	(B)	Zygomatic arch		(A)	Lingual nerve	
	(C)	Ramus of mandible		(B)	Chorda tympani	
	(D)	Carotid sheath		(C)	Nervous intermedius	
81.		lymphatic from the tip of the		(D)	Greater petrosal nerve	
	_	ue drains into :	86.	Drug	g causing hypocalcaemia is :	
	(A)	Submandibular lymph nodes		(A)	Lithium	
	(B)	Submental lymph nodes		(B)	Phenytoin	
	(C)	Jugulodigastric lymph nodes		(C)	Thiazides	
	(D)	Jugulo omohyoid lymph nodes		(D)	Vitamin A intoxication	
B – '	1C/34	(10	))		Contd.	

93.	by:  (A) Caffeine  (B) Amphetamine  (C) Cocaine  (D) Pethidine  Intolerance is:  (A) Toxic effect of a drug in an individual at therapeutic doses  (B) It indicates high threshold of the individual to the action of a drug	
93.	<ul> <li>(B) Amphetamine</li> <li>(C) Cocaine</li> <li>(D) Pethidine</li> <li>Intolerance is:</li> <li>(A) Toxic effect of a drug in an individual at therapeutic doses</li> <li>(B) It indicates high threshold of the individual to the action of a drug</li> </ul>	
93.	<ul> <li>(C) Cocaine</li> <li>(D) Pethidine</li> <li>Intolerance is:</li> <li>(A) Toxic effect of a drug in an individual at therapeutic doses</li> <li>(B) It indicates high threshold of the individual to the action of a drug</li> </ul>	
93.	<ul> <li>(D) Pethidine</li> <li>Intolerance is:</li> <li>(A) Toxic effect of a drug in an individual at therapeutic doses</li> <li>(B) It indicates high threshold of the individual to the action of a drug</li> </ul>	
93.	Intolerance is:  (A) Toxic effect of a drug in an individual at therapeutic doses  (B) It indicates high threshold of the individual to the action of a drug	
93.	<ul><li>(A) Toxic effect of a drug in an individual at therapeutic doses</li><li>(B) It indicates high threshold of the individual to the action of a drug</li></ul>	
	individual at therapeutic doses  (B) It indicates high threshold of the individual to the action of a drug	
	the individual to the action of a drug	
	drug	
	(C) Converse of the televerse	
	<ul><li>(C) Converse of the tolerance</li><li>(D) All of the above</li></ul>	
94.	Endocrinal side effects of psychotropic drugs are:	
	(A) Gynaecomastia	
	(B) Galactorrhea	
	(C) Both of the above	
	(D) None of the above	
95.	Drug of choice for febrile seizures :	
	(A) Diazepam	
	(B) Phenytoin	
	(C) Carbamate	
	(D) Valproate	
96.	Gynotype in a patient with Turner syndrome is:	
	(A) 45XO	
	(B) 47XXY	
	(C) 46XX	
	(D) 47XYY	
1)	(Turn over)	
	96.	

https://www.freshersnow.com/previous-year-question-papers

97.	·	most common site of intra-oral	102	. Soa	p bubble appearance is seen in :
	nevi	is:		(A)	Osteosarcoma
	(A)	Hard palate		(B)	Osteoclastoma
	(B)	Buccal muçosa		(C)	Ewing's sarcoma
	(C)	Gingiva		(D)	Multiple myeloma
	(D)	Lips	102	Deu	a used for the treatment of
98.		most common potentially gnant state of oral mucosa is:	103.		g used for the treatment of ostomia is:
	(A)	Leukoplakia		(A)	Atropine
	(B)	Erythroplakia		(B)	Ephidrine
• • .	(C)	Submucosal fibrosis		(C)	Scopolamine
	(D)	Keratoplakia		(D)	None of the above
99.		most common malignant plasm of the oral cavity is :	104.	Gag	reflex is lost due to the paralysis
	(A)	Squamous cell carcinoma		(A)	V <sup>th</sup> nerve
	(B)	Basal cell carcinoma		(B)	VII <sup>th</sup> nerve
	(C)	Malignant melanoma		(C)	IX <sup>th</sup> nerve
	(D)	Sarcoma		(D)	XII <sup>th</sup> nerve
100.	Unerrupted teeth is a common feature of :		105.		n's syndrome is trisomy of :
	(A)	Dentigerous cyst		(A)	16 chromosome
	(B)	Odontogenic keratocyst		(B)	18 chromosome
		Adenomatoid odontogenic		(C)	21 chromosome
		tumour		(D)	23 chromosome
	(D)	All of the above	106.	Pedo	odontics treatment triangle was
101.	Circu	moral pigmentation is seen in :		giver	ı by :
	(A)	Arsenic poisoning		(A)	G. Z. Wright
	(B)	Lead poisoning		(B)	Henry Kempe
	(C)	Peutz-Jaghers Syndrome		(C)	Evangeline Jordan
	(D)	Plummer-Vinson Syndrome		(D)	Addleston
B – 1	C/34	(12)	)		Contd,

107.		rior bite plane is used for the	112.		ch of the following GIC is using Pit and Fissure Sealent?
		Deep bite			
	(A) (B)	Crowding		(A)	Type 6
	(C)	Anterior cross bite		(B)	Type 8
	(D)	Posterior cross bite		(C)	Type 7
	• /			(D)	Type 9
108.		Daily dietary recommendation of	<sup>f</sup> 113.	The	first evidence of calcification of
		ium in children is :			nary teeth begins approximately
	(A)	600-800 mg		at:	
	(B)	800-1200 mg		(A)	11 weeks in utero
	(C)	1200-1600 mg		(B)	14 months in utero
	(D)	1600-2000 mg		(C)	16 months in utero
109.	Nick	el content in Stainless stee	I	(D)	6 months in utero
	crow	n is :	114.	Cla	ssical Conditioning was
	(A)	17-19%			posed by :
	(B)	10-13%		(A)	Skinner
	(C)	12-14%		(B)	Pavlov
	(D)	11-12%		(C)	Peaget
10.	Epst	ein Pearls are found :		(D)	Hassel
	(A)	On crest of ridges	4.48	, ,	
	(B)	Along mid palatine raphe	115.		sal Liability is :
	(C)	Midline of mandible		(A)	Maxilla-3mm, Mand-5mm
	(D)	On the buccal surface of incisor		(B)	Maxilla-7mm, Mand-5mm
111.	Perio	od of mixed dentition growth	ı	(C)	Maxilla-5mm, Mand-7mm
		t in boys :		(D)	Maxilla-5mm, Mand-3mm
	(A)	10-11 years	116.	ЕН	Angle introduced his classi-
	(B)	7-10 years		ficat	ion of malocclusion in :
	(C)	12-13 years		(A)	1988
	(D)	8-11 years		(B)	1899
B 1	1C/34	1	(13)	. ,	(Turn over)

https://www.freshersnow.com/previous-year-question-papers/

	(C)	1901		(C)	Buccinators
	(D)	1801		(D)	Temporalis
117.		most important synchondrosis for growth modification is :	122.		ine shift with unilateral posterior s bite may be due to a change
	(A)	Spheno ethmoidal		in:	
	(B)	Spheno occipital		(A)	Pitch
	(C)	Inter sphenoidal		(B)	Roll
	(D)	Inter maxillary		(C)	Yaw
118.	Mon	nent of a force is :		(D)	None of the above
	(A)	Force × perpendicular from centre of rotation	123.	Tent forth	h theorem of retention was put
	(B)	Force × perpendicular from centre of resistance		(A)	R. Riedel
	(C)	Area under the stress strain graph		(B) (C)	W. R. Proffit  A. Lundstrom
	(D)	Force × range		(D)	R. E. Moyers
119.	Third	l order bends are :	124.	Norn	nal naso labial angle necessi-
	(A)	In-out bends		tates	the extraction of :
	(B)	Tip back bends		(A)	Upper first premolar
	(C)	Torquing bends		(B)	Upper second premolar
	(D)	Anti-rotation bends		(C)	Upper incisor
120.	Ante	rior bolton's ratio is :		(D)	None of the above
	(A)	77.2%	125.	Reve	erse orthodontics refers to:
	(B)	91.3%		(A)	Repeated orthodontic treat-
	(C)	88.2%		( )	ment
	(D)	91.7%		(B)	Camouflage treatment
121.	Musc	cle affected in torticollis :		(C)	De-compensation
	(A)	Lateral pterygoid		(D)	Treatment using lingual
	(B)	Sternocleidomastoid			appliances
ر م <u>ا</u>	10121	(14)	,		Contd

126.	Junctional epithelium shows rete peg	g (C) Allograft
	formation in :	(D) Homograft
	(A) Stage I gingivitis	131. Ramping can be done for a :
	(B) Stage II gingivitis	(A) One wall defect
	(C) Stage III gingivitis	(B) Two wall defect
•	(D) Stage IV gingivitis	(C) Three wall defect
127.	Reunion of connective with root	t (D) Zero wall defect
	surface a root surface that has been pathologically exposed:	132. Halimeters are used in :
	(A) Reattachment	(A) Oral malodour
	(B) New attachment	(B) Tooth mobility
	(C) Regeneration	(C) Pathologic migration
	· · · · ·	(D) Pulp testing
128	(D) Repair  If half of the crown along with root is	133. Concept of full mouth disinfection
120.	separated or one half is removed, it	was put lotward by .
	is called :	(A) Miller
	(A) Radisection	(B) Glickman
	(B) Hemisection	(C) William
	(C) Apexification	(D) Quirynen
	(D) Coronosection	134. Epithelisation of gingivectomy wound is completed by:
129.	Most abundant immunoglobulin in the	
٠.	serum is:	(B) 1 month
	(A) IgG	(C) 4 months
	(B) IgA	(D) 2 months
:	(C) IgM	135. The microorganism found in higher
	(D) IgE	titers in pregnancy gingivitis :
130.	Which of the following grafts has the	(A) P. gingivalis
	best prognosis?	(B) P. intermedia
	(A) Xenograft	(C) Capnocytophaga
	(B) Autograft	(D) A. viscosus
IB – ′	1C/34	(15) (Turn over)

- 136. Each ml of 2% lignocaine contains:
  - (A) 0.2 mg of lignocaine
  - (B) 2 mg of lignocaine
  - (C) 10 mg of lignocaine
  - (D) 20 mg of lignocaine
- 137. Which of the following mandibular teeth may be removed without pain after the administration of IANB and lingual block?
  - (A) All molars
  - (B) Both premolars and molars
  - (C) All teeth in lower jaw
  - (D) First premolar and canine
- 138. Following structures are present in the submandibular space except:
  - (A) Deep part of submandibular gland
  - (B) Facial artery
  - (C) Mylohyoid nerve and vessels
  - (D) Superficial part of submandibular gland
- 139. Surgical emphysema may develop immediately after :
  - (A) Cutting the tooth with airotor
  - (B) Cutting the tooth with micromotor
  - (C) Cutting the bone with micromotor
  - (D) Cutting the bone with airotor

- 140. Metronidazole is a:
  - (A) Quinolone
  - (B) Anti-fungal
  - (C) Imidazole derivative
  - (D) Antiviral
- 141. Which of the following artery does not accompany the nerve throughout its course?
  - (A) Lingual
  - (B) Infra orbital
  - (C) Inferior alveolar
  - (D) Posterior superior alveolar
- 142. Secondary bony callus formation takes approximately:
  - (A) 1-2 weeks
  - (B) 3-4 weeks
  - (C) 4-6 weeks
  - (D) 6-8 weeks
- 143. Which of the following is DANGER SPACE?
  - (A) Pterygo mandibular
  - (B) Lateral pharyngeal
  - (C) Superficial and deep temporal
  - (D) Space between alar and prevertebral fascia
- 144. Weber Fergussion incision is used for:
  - (A) Mandibulectomy
  - (B) Maxillectomy
  - (C) Condylectomy
  - (D) Rhinoplasty

IB - 1C/34

(16)

145.	Cranial nerve passing through the centre of the cavernous sinus is:		150.	The target of gow gates technique is:		
	(A)	Optic nerve		(A)	Coronoid notch	
	(B)	Ophthalmic nerve		(B)	Sigmoid notch	
	(C)	Occulo motor nerve		(C)	Mandibular foramen	
	(D)	Abducent nerve		(D)	Neck of condyle	
146.	Artic	ular disc of TMJ is :	151.	"Ent	er bolus-exit balance" was stated	
	(A)	Fibro cartilage		by:	• • •	
	(B)	Hyaline cartilage		(A)	Harold Ortman	
	(C)	Elastic tissue		(B)	Sheppard	
	(D)	TMJ ligament		(C)	Stewart	
447		•		(D)	Hudson	
147.	Four clotting factors synthesized in liver are:		152.		estimated width of maxillary	
·. ·	(A)	III, V, VII and VIII			ral incisor is:	
	(B)	II, VII, IX and X		(A)	1/16 <sup>th</sup> bizygomatic width	
	(C)	II, VI, IX and XII		(B)	1/16 <sup>th</sup> interpupillary distance	
	(D)	II, V, VII and IX		(C)	1/8 <sup>th</sup> bizygomatic width	
440				(D)	None of the above	
140.	Which of the following is efficacious modality in diagnosis of salivary gland neoplasm?		153.	In a	In arcon articulators the condylar element is located in :	
				elen		
	(A)			(A)	Upper member	
	(B)	Excision biopsy		(B)	Lower member	
	(C)	Fine needle aspiration cytology		(C)	No condyle	
	(D)	MRI scan		(D)	Fixed condyle	
			154.	The	quality that distinguishes one	
149.	1 : 1000 adrenaline ampule contains :			from the other is :		
	(A)	0.1 mg/ml		(A)	Hue	
	(B)	1 mg/ml		(B)	Chroma	
	(C)	10 mg/ml		(C)	Value	
	(D)	0.01 mg/ml		(D)	Saturation	

(17)

IB - 1C/34

(Turn over)

155,	In laminate technique of impression making:			Instrument used to locate pterygomaxillary notch:	
	(A)	Alginate and agar are used		(A)	Boby gauge
	(B)	Only alginate is used		(B)	Divider
	(C)	Elastomers are used		(C)	Screw gauge
	(D)	Only agar		(D)	T-burnisher
4 d S	ante papp (A) (B) (C) (D)	tral incisors are generally rior to the line bisecting incisive bilae by:  8-10 mm  14-15 mm  2-6 mm  11-20 mm  antages of an over denture:  Preservation of ridge		The dent (A) (B) (C)	distal border of mandibular ture should cover:  Ant. $\frac{1}{3}$ rd of retromolar pad  Ant. $\frac{2}{3}$ rd of retromolar pad  Should not cover the retromolar pad  Hamullar notch  rylic resin, hydroquinone act as:  Activator
. /	(B)			(B)	Initiator
	` ,	Retention		(C)	Inhibitor
	(D)	,		(D)	Plasticizer
158.	(A) (B) (C) (D)	bow transfer records : Orientation relation Special relation Vertical relation Tentative centric relation	163.		nate is made dust free by the tion of : Glycerine Calcium sulphate Sodium phosphate
159.	Ident	ntify the correct hanau formula	- ' '	(D)	Calcium chloride
	which gives the lateral condylar		404	` ,	
	guidance angle :		164.		seat can be prepared:
	(A)	L = H/8 + 12		(A)	Only on tooth surface
	` '	L = 8/H + 12		(B)	In amalgam restoration
	(C)	L = H/8 * 12		(C)	In cast gold restoration
	(D)	L = H/8 – 12		(D)	All of the above

(18)

IB - 1C/34

165	. Hea	l raising movement of a RPD	is		(C)	Histogram	
	prevented by:				(D)	Pie diagram	
	(A)	Occlusal rest	1	70	Δn	experiment in which the	
	(B)	Minor connector	•	70.		perimental subjects cannot	
	(C)	Direct retainer				tinguish the experimental	
	(D)	Indirect retainer				nditions is called as :	
166.	Indi	cator in Snyder Test is :	٠		(A)	Two Tailed Testing	
	(A)	Diazoresorcinol			(B)	Single Blind experiment	
	(B)	Methyl red			(C)	Double Blind experiment	
	(C)	Glucose			(D)	Retrospective experiment	
	(D)	Bromocresol green	1	71.	Pit a	and fissure sealant is a form of :	
167.	First	artificial water fluoridation of	of		(A)	Primary prevention	
	wate	water supply was done in :			(B)	Secondary prevention	
	(A)	London			(C)		
,	(B)	New York			• ,	Tertiary prevention	
	(C)	Grand Rapids			(D)	Primordial prevention	
	(D)	Ontario, Canada	• 17	72.	Sho	e leather survey was carried out	
168.	Specificity of a test denotes :				by:		
	(A)	) True Positive			(A)	Churchill H. V.	
	(B)	True Negative			(B)	Frederick McKay	
	(C)	False Positive			(C)	Trendley H. Dean	
	(D)	False Negative			(D)	Greene Vardiman Black	
169.	The coverage of the geographic area through dental clinics can be		17	<b>7</b> 3.	Exar	mple of a composite index is :	
					(A)	CPITN	
	depid	cted by the following way :			(B)	DMF	
	(A)	Frequency polygon			(C)	TNM	
	(B)	Cartogram			(D)	Periodontal	
IB – 1C/34 (		(19)			(Turn over)		

https://www.freshersnow.com/previous-year-question-papers/

174.	-	rrence of disease clearly in		(B)	Permanent mandibular canine	
	excess of expected occurrence:			(C)	Deciduous mandibular canine	
	(A)	Endemic		(D)	Maxillary second premolar	
	(B)	(B) Pandemic		Shar	pe of occlusal aspect of maxillary	
	(C)	Sporadic	170.	permanent first molar is:		
	(D)	Epidemic		(A)	Trapezoidal	
175.	Which of the following is a measure			(B)	Rectangle	
	of dispersion?			` '	_	
	(A)	Mean		(C)	Square	
	(B)	Median		(D)	Rhomboid	
	(C)	Mode	180.	Whi	ch of the following is not a groove	
	(D)	Standard deviation		patt	pattern of mandibular second	
176.	Which of the following deciduous			premolar?		
170.	tooth does not resemble any of the			(A)	Υ	
	tooth in permanent dentition?			(B)	Н	
	(A)	<b>A</b> · .		(C)	U	
	(B)	В	181.	(D)	M	
	(C)	J		Cell	s concerned with resorption of	
	(D)	L .	, , , , ,		deciduous teeth:	
177.	Which of the following permanent tooth has a single antagonist in occlusion?			(A)	Osteoblast	
				(B)	Cementoclast	
				(C)	Odontoclast	
	(A)	Maxillary third molar		(D)	Pericytes	
	(B)	Mandibular third molar		• •	•	
	(C)	Mandibular canine	182.		ch of the following tooth contain	
	(D)	Maxillary central incisor			olementary grooves ?	
178.	Whi		(A)	Maxillary second premolar		
	mesial cusp slope greater than distal			(B)	Mandibular second molar	
	cusp slope?			(C)	Both (A) and (B)	
	(A)	Maxillary first premolar		(D)	None of the above	

		·			
183	. Col	is seen in :	188	3. Ack	terman's tumor is :
	(A)	Buccal mucosa		(A)	Pleomorphic adenoma
•	(B)	Inter dental papilla		(B)	Verrucous carcinoma
	(C)	Floor of mouth		(C)	Spindle cell carcinoma
	(D)	Hard palate		(D)	Adenolymphoma
184	. Epithelial rests of Malassez is seen in :		189		ich of the following salivary gland or is mimic mucocele?
	(A)	Palate		(A)	Adenoid cystic carcinoma
•	(B)	Gingiva		(B)	Salivary duct carcinoma
	(C)	Periodontal ligament		(C)	Mucoepidermoid carcinoma
	(D)	Dental pulp		(D)	Sebaceous carcinoma
195			190	. Odc	ontoclastoma is :
100,	Mesiolingual groove is seen in which of the following teeth?			(A)	Giant cell tumor of bone
	(A)	Maxillary lateral incisor		(B)	Giant cell granuloma
	(B)	•		(C)	Internal resorption of tooth
	` .	Mandibular first premolar		(D)	Hypercementosis
	(C)	Permanent mandibular firşt molar	191.	Mun	nps is caused by :
	(D)	Mandibular canine		(A)	Orthomyxovirus
400				(B)	Adenovirus
186.		ng's Premolar is :		(C)	Paramyxovirus
	(A)	Dens invaginatus		(D)	Coxsackie virus
	(B)	Dens evaginatus		Odo	ntoclasia is : an igraphy and a second
	(C)	Taurodontism		(A)	Linear enamel caries
	(D)	Enamel hypoplasia		(B)	Tooth resorption
187.	Most	common benign tumor of oral		(C)	Odontogenic tumor
	cavity is :		•	(D)	Dry socket
	(A)	Papilloma	193	Bifid	rib is seen in which of the
	(B)	Nevus	.00.		ving syndrome?
	(C)	Fibroma		(A)	Papillon Leferve syndrome
	(D)	Lipoma		(B)	Gorlin syndrome

(21)

IB-1C/34

(Turn over)

- (C) Sturge Weber syndrome (D) Ascher syndrome 194. Most common oral manifestation of diabetes mellitus is: (A) Leukoplakia (B) Periodontal abscess (C) **ANUG** (D) Hyperplastic candidiasis 195. Alarm clock headache is: Trigeminal neuralgia (A) Temporal arteritis (B)
- 196. Drug of choice for the treatment of Actinomycosis is:

Sphenopalatine neuralgia

Glossopharyngeal neuralgia

(A) Penicillin

(C)

- (B) Cephalosporin
- (C) Clotrimazole
- (D) Metronidazole
- 197. Moeller's Glossities is seen in :
  - (A) Iron deficiency Anaemia

- (B) Vitamin B6 deficiency
- (C) Vitamin B12 deficiency
- (D) Thiamin deficiency
- 198. Life span of platelet is:
  - (A) 6-8 days
  - (B) 12-14 days
  - (C) 60 days
  - (D) 120 days
- 199. Rushton bodies are present in:
  - (A) Periodontal cyst
  - (B) Dentigerous cyst
  - (C) Odontogenic keratocyst
  - (D) Stafne's cyst
- 200. Muscle which depresses the mandible is:
  - (A) Lateral Pterygoid
  - (B) Medial Pterygoid
  - (C) Temporalis
  - (D) Masseter

## SPACE FOR ROUGH WORK

B

 $C_{\ell}$