

Harmonic Oscillator

For a Harmonic Oscillator potential energy $V = \frac{1}{2} kx^2$, k is the force constant, is a continuous function of the coordinate x and therefore is completely different from systems we considered so far where the potential is constant over a region.

Wave Equation

$$\begin{aligned} V &= \frac{1}{2} kx^2 \\ &= \frac{1}{2} 4\pi m V_0^2 x^2 \\ &= \frac{1}{2} m\omega^2 x^2 \end{aligned}$$

The time - independent Schrödinger equation of the linear harmonic oscillator then

$$\frac{\partial^2 \Psi}{\partial x^2} + \frac{2m}{\hbar^2} (E - \frac{1}{2} m\omega^2 x^2) \Psi = 0 \rightarrow (1)$$

It is convenient to work with a new variable y and a new parameter λ by

$$y = \left(\frac{m\omega}{\hbar}\right)^{1/2} x \quad \text{and} \quad \lambda = \frac{2E}{\hbar\omega}$$

In terms of these quantities schrödinger equation can be written as

$$\frac{\partial^2 \Psi}{\partial x^2} + (\lambda - y^2) \Psi = 0 \rightarrow (2)$$

Asymptotic solution

We have to find the solution when $y \rightarrow \infty$, when y is very large, $\lambda - y^2 \cong -y^2$

$$\text{Equation (2) becomes } \frac{\partial^2 \Psi}{\partial x^2} - y^2 \Psi = 0 \rightarrow (3)$$

Its asymptotic solutions are

$$\begin{aligned} \Psi(x) &= e^{\pm y^2/2} \\ (2) \rightarrow \frac{\partial^2 \Psi}{\partial x^2} &= (y^2 + 1) e^{\pm y^2/2} \end{aligned}$$

Out of the two asymptotic solutions $e^{y^2/2}$ is not acceptable as it diverges

When $|y| \rightarrow \infty$. The exact solution can be written as

$$\Psi = e^{-y^2/2} H(y)$$

Where $H(y)$ is a function of y and the product $e^{-y^2/2} H(y)$ tends to Zero as

$$|y| \rightarrow \infty$$

Series Solution

Substituting the value of Ψ in (3)

$$(3) \rightarrow \frac{d^2 H(y)}{dy^2} - 2y \frac{dH}{dy} + (\lambda - 1) H = 0$$

Which is known as the Hermite equation. We shall look for $H(y)$, a series solution of the type.

$$H(y) = \sum_{n=0}^{\infty} a_n y^n \rightarrow (4)$$

Substituting the value of $H(y)$ in equation (3)

$$\sum_{k=0}^{\infty} [(k+1)(k+2)a_{k+2} - (2k+1-\lambda)a_k]y^k = 0 \rightarrow (5)$$

For the validity of this equation, coefficient of each power of y must vanish separately. Coefficient of y^k when equated to zero gives the recurrence relation.

$$a_{k+2} = \frac{2k+1-\lambda}{(k+1)(k+2)} \rightarrow (6)$$

Equation (4) will have only odd coefficients if $a_0=0$ and even coefficients if $a_1 = 0$.

Thus we have two independent solutions for $H(y)$ and a linear combination of the two will be the most general solution. The two solutions are ;

$$H_e(y) = a_0 + a_2 y^2 + a_4 y^4 + \dots$$

$$H_o(y) = y (a_1 + a_3 y^2 + a_5 y^4 + \dots)$$

Energy Eigenvalues

When $K \rightarrow \infty$ we get

$$\frac{a_{k+2}}{a_k} = \frac{2}{k}$$

Consider the Taylor series expansion of $\exp(y^2)$

$$\exp(y^2) = \sum_{0,2,4} \frac{1}{(k/2)!} y^k \rightarrow (7)$$

The ratio of coefficients of the successive terms in equation (7) is

$$\frac{a_{k+2}}{a_k} = \frac{(k/2)!}{[(k/2)+1]!} = \frac{1}{(k/2)+1} = \frac{2}{k}$$

Where k is large. Therefore, for large values of k , $\Psi = \exp\left(\frac{y^2}{2}\right)$ tends to behave like $\exp\left(\frac{y^2}{2}\right)$, if the series is even, and $y \exp\left(\frac{y^2}{2}\right)$ if this series is odd; which is not acceptable. Substitution of λ gives

$$2n+1 - \frac{2E}{\hbar\omega} = 0$$

$$E_n = \left(n + \frac{1}{2}\right)\hbar\omega \quad n = 0, 1, 2, \dots$$

The energy of an oscillator based on quantum theory is

$$E_n = n\hbar\omega$$

From the two expressions, it is evident that the quantum mechanical energy value is higher than the quantum theory value by $\frac{1}{2}\hbar\omega$, which is the energy possessed by the lowest state $n = 0$. The oscillator will have this energy even at absolute zero. This energy of $\frac{1}{2}\hbar\omega$ is called the zero-point energy which is a manifestation of uncertainty principle. $E=0$ means that both position and momentum are well defined which is a violation of uncertainty principle. It can be shown that the minimum energy of an oscillator without violating uncertainty principle $\frac{1}{2}\hbar\omega$.

Energy Eigenfunctions

When $\lambda = 2n + 1$ Eq (3) reduces to

$$\frac{d^2 H_n(y)}{dy^2} - 2y \frac{dH_n(y)}{dy} + 2nH_n(y) = 0$$

The energy eigenfunctions can be expressed as

$$\Psi_n(y) = N_n H_n(y) \exp\left(-\frac{y^2}{2}\right), \quad y^2 = \frac{m\omega}{\hbar} x^2$$

The first - four Hermite polynomials are

$$H_0(y) = 1$$

$$H_1(y) = 2y$$

$$H_2(y) = 4y^2 - 2$$

$$H_3(y) = 8y^3 - 12y$$

The normalization condition leads to

$$|N_n|^2 \left(\frac{\hbar}{m\omega}\right)^{1/2} \int_{-\infty}^{\infty} H_n^2(y) \exp(-y^2) dy = 1$$

The normalized eigenfunctions and ground state eigenfunctions are given by

$$\Psi_n(y) = \left[\frac{(m\omega)^{1/2}}{(h\pi)} \frac{1}{2^n(n)!} \right]^{1/2} H_n(y) \exp\left(-\frac{y^2}{2}\right)$$

$$\Psi_0(x) = \left(\frac{m\omega}{h\pi}\right)^{1/4} \exp\left(-\frac{m\omega x^2}{2\hbar}\right)$$

The wave functions $\Psi_n(x)$ and the probability density $|\Psi_n(x)|^2$ of the lowest four states are illustrated below.




Figure ; (a) Energy levels (E_n) and wave functions $\Psi_n(x)$ of the lowest four states of the linear harmonic oscillator. (b) probability density $|\Psi_n(x)|^2$ of the lowest four states.