

NUCLEAR FORCES

- The protons and neutrons are held together by the strong attractive forces inside the nucleus, These forces are called as nuclear forces.

properties of the Nuclear Force

- Nuclear force is short ranged. It exists in small region (of diameter $m = 1 \text{ fm}$). The nuclear force between two nucleons decreases rapidly as the separation between them increases and becomes negligible at separation more than 10

fm.

- Nuclear force is much stronger than electromagnetic force and gravitational force.
- Nuclear force is independent of charge. The nuclear force between two protons is same as that between two neutrons or between a neutron and proton. This is known as charge independent character of nuclear force.

Nuclear Reaction


- In nuclear reaction, sum of masses before reaction is greater than the sum of masses after the reaction. The difference in masses appears in the form of energy following the law of inter-conversion of mass and energy. The energy released in a nuclear reaction is called as Q value of a reaction and is given as follows .
 - If difference in mass before and after the reaction is Δm
- u

$\Delta m = \text{mass of reactants} - \text{mass of products}$, then Q value = $\Delta m(931) \text{ MeV}$

- Law of conservation of momentum is also followed.
- Total number of protons and neutrons should also remain same on both sides of a nuclear reaction.

Nuclear Fission

- The breaking of a heavy nucleus into two or more fragments of comparable masses, with the release of tremendous amount of energy is called as nuclear fission. The most typical fission reaction occurs when slow moving neutrons strike ${}_{92}\text{U}^{235}$. The following nuclear reaction takes place.




- If more than one of the neutrons produced in the above fission reaction are capable of inducing a fission reaction (provided U^{235} is available), then the number of fissions taking place at successive stages goes increasing at a very brisk rate and this generates a series of fission reactions. This is known as chain reaction. If mass of U^{235} sample greater than a certain size called the critical size then it is capable of continuous fission by itself.
- If the number of fissions in a given interval of time goes on increasing continuously, then a condition of explosion is created. In such cases, the chain reaction is known as uncontrolled chain reaction. This forms the basis of atomic bomb.

- In a chain reaction, the fast moving neutrons are absorbed by certain substances known as moderators (like heavy water), then the number of fissions can be controlled and the chain reaction in such case is known as controlled chain reaction. This forms the basis of a nuclear reactor.

Nuclear Fusion

- The process in which two or more light nuclei are combined into a single nucleus with the release of tremendous amount of energy is called as nuclear fusion. Like a fission reaction, the sum of masses before the fusion (i.e. of light nuclei) is more than the sum of masses after the fusion (i.e. of bigger nucleus) and this difference appears as the fusion energy. The most typical fusion reaction is the fusion of two deuterium nuclei into helium.



- For the fusion reaction to occur, the light nuclei are brought closer to each other (with a distance of 10^{-14} m). This is possible only at very high temperature to counter the repulsive force between nuclei. Due to this reason, the fusion reaction is very difficult to perform, The inner core of sun is at very high temperature, and is suitable for fusion, in fact the source

of energy in sun and other stars is the nuclear fusion reaction.