

## 15. Quark Model

### 15.1 Quantum numbers of the quarks

Quantum chromodynamics (QCD) is the theory of strong interactions. QCD is a quantum field theory and its constituents are a set of fermions, the quarks, and gauge bosons, the gluons. Strongly interacting particles, the hadrons, are bound states of quark and gluon fields. As gluons carry no intrinsic quantum numbers beyond color charge, and because color is believed to be permanently confined, most of the quantum numbers of strongly interacting particles are given by the quantum numbers of their constituent quarks and antiquarks. The description of hadronic properties which strongly emphasizes the role of the minimum-quark-content part of the wave function of a hadron is generically called the quark model. It exists on many levels: from the simple, almost dynamics-free picture of strongly interacting particles as bound states of quarks and antiquarks, to more detailed descriptions of dynamics, either through models or directly from QCD itself. The different sections of this review survey the many approaches to the spectroscopy of strongly interacting particles which fall under the umbrella of the quark model.

Quarks are strongly interacting fermions with spin  $1/2$  and, by convention, positive parity. Antiquarks have negative parity. Quarks have the additive baryon number  $1/3$ , antiquarks  $-1/3$ . Table 15.1 gives the other additive quantum numbers (flavors) for the three generations of quarks. They are related to the charge  $Q$  (in units of the elementary charge  $e$ ) through the generalized Gell-Mann-Nishijima formula

$$Q = I_z + \frac{B + S + C + B + T}{2}, \quad (15.1)$$

where  $B$  is the baryon number. The convention is that the quark *flavor* ( $I_z$ ,  $S$ ,  $C$ ,  $B$ , or  $T$ ) has the same sign as its *charge*  $Q$ . With this convention, any flavor carried by a charged meson

**Table 15.1**

	$d$	$u$	$s$	$c$	$b$	$t$
Q – electric charge	$-\frac{1}{3}$	$+\frac{2}{3}$	$-\frac{1}{3}$	$+\frac{2}{3}$	$-\frac{1}{3}$	$+\frac{2}{3}$
$I$ – isospin	$\frac{1}{2}$	$\frac{2}{3}$	$0$	$0$	$0$	$0$
$I_z$ – isospin z-component	$-\frac{1}{2}$	$+\frac{1}{2}$	$0$	$0$	$0$	$0$
$S$ – strangeness	$0$	$0$	$-1$	$0$	$0$	$0$
$C$ – charm	$0$	$0$	$0$	$+1$	$0$	$0$
$B$ – bottomness	$0$	$0$	$0$	$0$	$-1$	$0$
$T$ – topness	$0$	$0$	$0$	$0$	$0$	$+1$

has the same sign as its charge, *e.g.*, the strangeness of the  $K^+$  is  $+1$ , the bottomness of the  $B^+$  is  $+1$ , and the charm and strangeness of the  $D_s^-$  are each  $-1$ . Antiquarks have the opposite flavor signs. The hypercharge is defined as

$$Y = B + S - \frac{C - B + T}{3}. \quad (15.2)$$

Thus  $Y$  is equal to  $\frac{1}{3}$  for the  $u$  and  $d$  quarks,  $-\frac{2}{3}$  for the  $s$  quark, and  $0$  for all other quarks. More details and derivations on the quark structure of mesons and baryons can be found in Ref. [1].

## 15.2 Mesons

Mesons have baryon number  $B = 0$ . In the quark model, they are  $qq^{-0}$  bound states of quarks  $q$  and antiquarks  $q^{-0}$  (the flavors of  $q$  and  $q^{-0}$  may be different). If the orbital angular momentum of the  $qq^{-0}$  state is  $\ell$ , then the parity  $P$  is  $(-1)^{\ell+1}$ . The meson spin  $J$  is given by the usual relation  $|\ell - s| \leq J \leq \ell + s$ , where  $s$  is 0 (antiparallel quark spins) or 1 (parallel quark spins). The charge conjugation, or  $C$ -parity  $C = (-1)^{\ell+s}$ , is defined only for the  $qq^{-}$  states made of quarks and their own antiquarks. The  $C$ -parity can be generalized to the  $G$ -parity  $G = (-1)^{\ell+s}$  for mesons made of quarks and their own antiquarks (isospin  $I_z = 0$ ), and for the charged  $ud^{-}$  and  $du^{-}$  states (isospin  $I = 1$ ).

The mesons are classified in  $J^{PC}$  multiplets. The  $\ell = 0$  states are the pseudoscalars ( $0^{-+}$ ) and the vectors ( $1^{-}$ ). The orbital excitations  $\ell = 1$  are the scalars ( $0^{++}$ ), the axial vectors ( $1^{++}$ ) and ( $1^{+-}$ ), and the tensors ( $2^{++}$ ). Assignments for many of the known mesons are given in Tables 15.2, 15.3 and 15.4. Radial excitations are denoted by the principal quantum number  $n$ . The very short lifetime of the  $t$  quark makes it likely that bound-state hadrons containing  $t$  quarks and/or antiquarks do not exist.

States in the natural spin-parity series  $P = (-1)^J$  must, according to the above, have  $s = 1$  and hence,  $CP = +1$ . Thus, mesons with natural spin-parity and  $CP = -1$  ( $0^{+-}$ ,  $1^{+-}$ ,  $2^{+-}$ ,  $3^{+-}$ , etc.) are forbidden in the  $qq^{-0}$  model. The  $J^{PC} = 0^{-}$  state is forbidden as well. Mesons with such *exotic* quantum numbers may exist, but would lie outside the  $qq^{-0}$  model (see section below on exotic mesons).

Following  $SU(3)$ , the nine possible  $qq^{-0}$  combinations containing the light  $u$ ,  $d$ , and  $s$  quarks are grouped into an octet and a singlet of light quark mesons:

$$\mathbf{3} \otimes \mathbf{\bar{3}} = \mathbf{8} \oplus \mathbf{1}. \quad (15.3)$$

A fourth quark such as charm  $c$  can be included by extending  $SU(3)$  to  $SU(4)$ . However,  $SU(4)$  is badly broken owing to the much heavier  $c$  quark. Nevertheless, in an  $SU(4)$  classification, the sixteen mesons are grouped into a 15-plet and a singlet:

$$\mathbf{4} \otimes \mathbf{\bar{4}} = \mathbf{15} \oplus \mathbf{1}. \quad (15.4)$$

The *weight diagrams* for the ground-state pseudoscalar ( $0^{-+}$ ) and vector ( $1^{-}$ ) mesons are depicted in Fig. 15.1. The light quark mesons are members of nonets building the middle plane in Fig. 15.1(a) and (b).

Isoscalar states with the same  $J^{PC}$  mix, but mixing between the two light quark isoscalar mesons, and the much heavier charmonium or bottomonium states, are generally assumed to be negligible. In the following, we shall use the generic names  $a$  for the  $I = 1$ ,  $K$  for the  $I = 1/2$ , and  $f$  and  $f'$  for the  $I = 0$  members of the light quark nonets. Thus, the physical isoscalars are mixtures of the  $SU(3)$  wave function  $\psi_8$  and  $\psi_1$ :

$$f' = \psi_8 \cos\vartheta - \psi_1 \sin\vartheta, \quad (15.5)$$

$$f = \psi_8 \sin\vartheta + \psi_1 \cos\vartheta, \quad (15.6)$$

where  $\vartheta$  is the nonet mixing angle and

$$\psi_8 = \frac{1}{\sqrt{6}}(u\bar{u} + d\bar{d} - 2s\bar{s}), \quad (15.7)$$

$$\psi_1 = \frac{1}{\sqrt{3}}(u\bar{u} + d\bar{d} + s\bar{s}). \quad (15.8)$$

**Table 15.2:** Suggested  $qq^-$  quark-model assignments for some of the observed light mesons. Mesons in bold face are included in the Meson Summary Table. The wave functions  $f$  and  $f^0$  are given in the text (Eqn. 15.9). The singlet-octet mixing angles from the linear mass formula (15.12) and its quadratic version (in which the masses are squared) are also given for the well established nonets. The classification of the  $0^{++}$  mesons is tentative: the light scalars  $a_0(980)$ ,  $f_0(980)$ ,  $f_0(500)$  and  $K_0^*$  (700) are often considered to be four-quark states, and are omitted from the table, see Eqn. (15.26) below. The isoscalar  $0^{++}$  mesons  $f_0(1370)$ ,  $f_0(1500)$  (not shown) and  $f_0(1710)$  are expected to mix, see the “Note on Non- $qq^-$  mesons” and the “Note on Scalar Mesons below 2 GeV” in the Meson Listings for details. The isoscalar assignments in the  $2^1S_0$  ( $0^+$ ) nonet are also tentative. The  $\eta(1405)$  (not shown) and  $\eta(1475)$  may be manifestations of the same state, see the “Note on Pseudoscalar and Pseudovector Mesons in the 1400 MeV Region” in the Meson Listings.

<sup>†</sup> The  $1^{++}$  and  $2^{--}$  isospin  $\frac{1}{2}$  states mix. In particular, the  $K_{1A}$  and  $K_{1B}$  are nearly equal ( $45^\circ$ ) mixtures of the  $K_1(1270)$  and  $K_1(1400)$  (see [2] and references therein). <sup>‡</sup> The physical vector mesons may be mixtures of  $1^3D_1$  and  $2^3S_1$  [3].

$n2s+1 \backslash JPC \quad I = 1$	$I = \frac{1}{2}$	$I = 0$	$I = 0$	$\vartheta_{\text{quad}}$	$\vartheta_{\text{lin}}$
$J$				$^\circ$	$^\circ$
$ud^-, ud^-,$ $\frac{1}{\sqrt{2}}(d\bar{d} - u\bar{u})$	$us^-, ds^-; us$ $\bar{d}s, -$	$f_0$	$f$		
<b>11S 0- <math>\pi</math></b> 0 +	<b><math>K</math></b>	<b><math>\eta</math></b>	<b><math>\eta^0(958)</math></b>	-11.3	-24.5
<b>13S 1- <math>\rho(770)</math></b> 1 -	<b><math>K^*(892)</math></b>	<b><math>\phi(1020)</math></b>	<b><math>\omega(782)</math></b>	39.	36.
<b>11P 1+ <math>b_1(1235)</math></b> 1 -	<b><math>K_{1B}^+</math></b>	<b><math>h_1(1415)</math></b>	<b><math>h_1(1170)</math></b>	2	5
0+ $a$ +	$K_0^*(1430)$	$f_0(1710)$	$f_0(1370)$		
<b>13P 1+ <math>a_1(1260)</math></b> 1 +	<b><math>K_{1A}^+</math></b>	<b><math>f_1(1420)</math></b>	<b><math>f_1(1285)</math></b>		
2+ $a$ +	$K_2^*(1430)$	$f_2(1270)$	$f_2(1270)$	29.	28.
<b>11D 2- <math>\pi_2(1670)</math></b> 2 +	<b><math>K_2(1770)^+</math></b>	<b><math>\eta_2(1870)</math></b>	<b><math>\eta_2(1645)</math></b>	6	0
<b>13D 1- <math>\rho(1700)</math></b> 1 -	<b><math>K^*(1680)^+</math></b>	$\omega(1650)$			
<b>13D 2-</b> 2 -	<b><math>K_2(1820)^+</math></b>				
3- <b><math>\rho_3(1690)</math></b> -	<b><math>K_3^*(1780)</math></b>	$\phi_3(1850)$	$\omega_3(1670)$	31.	30.
4+ $a$ +	$K_4^*(2040)$	$f_4(2300)$	$f_4(2050)$	8	8
5- -	$K_5^*(2380)$				
<b>21S 0- <math>\pi(1300)</math></b> 0 +	<b><math>K(1460)</math></b>	<b><math>\eta(1475)</math></b>	<b><math>\eta(1295)</math></b>		

$2_{3S} 1^- \rho(1450)$	$K^*(1410)^{\pm}$	$\varphi(1680) \omega(1420)$
$1^-$		$)$
$2_{3P} 1^+ a_1(1640)$		
$1^+$		
$2^+ a$	$K_2^*(1980)$	$f_2(1950) f_2(1640)$
$+$		

These mixing relations are often rewritten to exhibit the  $uu\bar{u} + dd\bar{d}$  and  $ss\bar{s}$  components which decouple

for the “ideal” mixing angle  $\vartheta_i$ , such that  $\tan \vartheta_i = 1/\sqrt{2}$  (or  $\vartheta_i = 35.3^\circ$ ). Defining  $\alpha = \vartheta + 54.7^\circ$ , one obtains the physical isoscalar in the flavor basis

$$f' = \frac{1}{\sqrt{2}}(u\bar{u} + d\bar{d}) \cos \alpha - s\bar{s} \sin \alpha, \quad (15.9)$$

and its orthogonal partner  $f$  (replace  $\alpha$  by  $\alpha - 90^\circ$ ). Thus for ideal mixing ( $\alpha_i = 90^\circ$ ), the  $f^0$  becomes pure  $ss\bar{s}$  and the  $f$  pure  $uu\bar{u} + dd\bar{d}$ . The mixing angle  $\vartheta$  can be derived by diagonalizing the mass matrix

$$\begin{pmatrix} m_8 & m_{81} \\ m_{18} & m_1 \end{pmatrix} \quad (15.10)$$

**Table 15.3:**  $cc\bar{c}$  quark-model assignments for the charmonium and open charm mesons with established  $J^{PC}$ . Mesons in bold face are included in the Meson Summary Table. The open flavor states in the  $1^{+-}$  and  $1^{++}$  rows are mixtures of the  $1^{+\pm}$  states.

<sup>†</sup>The masses are considerably smaller than most theoretical predictions.

These states have also been considered as four-quark states. <sup>#</sup>  
Mixtures of the  $1^3D_1$  and  $2^3S_1$  states.

$n2s+1J$	$J^{PC}$	$I = 0$	$I = \frac{1}{2}$	$I = 0$
		$cc\bar{c}$	$cu\bar{c}, cd\bar{c};$ $cu\bar{u}, cd\bar{d}$	$cs\bar{c};$ $cs\bar{s}$
$1^1S_0$	$0^{-+}$	<b><math>\eta_c(1S)</math></b>	<b><math>D</math></b>	<b><math>D_{s\pm}</math></b>
$1^3S_1$	$1^{--}$	<b><math>J/\psi(1S)</math></b>	<b><math>D^*</math></b>	<b><math>D_{s^* \pm}</math></b>
$1^3P_0$	$0^{++}$	<b><math>\chi_{c0}(1P)</math></b>	<b><math>D_0^*(2300)</math></b>	<b><math>D_{s0}^*(2317)^{\pm \dagger}</math></b>
$1^3P_1$	$1^{++}$	<b><math>\chi_{c1}(1P)</math></b>	$D_1(2430)$	<b><math>D_{s1}(2460)^{\pm \dagger}</math></b>
$1^1P_1$	$1^{+-}$	<b><math>h_c(1P)</math></b>	<b><math>D_1(2420)</math></b>	<b><math>D_{s1}(2536)^{\pm \dagger}</math></b>
$1^3P_2$	$2^{++}$	<b><math>\chi_{c2}(1P)</math></b>	<b><math>D_2^*(2460)</math></b>	<b><math>D_{s2}^*(2573)</math></b>
$2^1S_0$	$0^{-+}$	<b><math>\eta_c(2S)</math></b>		
$2^3S_1$	$1^{--}$	<b><math>\psi(2S)</math></b>		<b><math>D_{s^*1}(2700)^{\pm \dagger}</math></b>
$1^3D_1$	$1^{--}$	<b><math>\psi(3770)</math></b>		<b><math>D_{s1}^*(2860)^{\pm \dagger}</math></b>
$1^3D_2$	$2^{--}$	<b><math>\psi_2(3823)</math></b>		
$2^3P_J$	$0, 1^{++}$	$\chi_{c0}(3860)$		

	2 <sup>++</sup>	$\chi_{c2}(3930)$	
3 $3S_1$	1 <sup>--</sup>	$\psi(4040)$	
2 $3D_1$	1 <sup>--</sup>	$\psi(4160)$	
4 $3S_1$	1 <sup>--</sup>	$\psi(4415)$	
1 $3D_3$	3 <sup>--</sup>	$D_3^*(2750)$	$D_{s3}^*(2860)^\pm$

The mass eigenvalues are  $m_{f0}$  and  $m_f$ . The mixing angle is given by

$$\tan\vartheta = \frac{m_8 - m_{f0}}{m_{81}}. \quad (15.11)$$

Calculating  $m_8$  and  $m_{81}$  from the wave functions Eq. 15.7 and Eq. 15.8, and expressing the quark masses as a function of the  $I = 1/2$  and  $I = 1$  meson masses, one obtains

$$\tan\vartheta = \frac{4m_K - m_a - 3m_{f0}}{2(m_a - m_K)} \sqrt{2}, \quad (15.12)$$

which also determines the sign of  $\vartheta$ . Alternatively, one can express the mixing angle as a function of all nonet masses. The octet mass is given by

$$m_8 = m_{f0} \cos^2 \vartheta + m_f \sin^2 \vartheta \quad (15.13)$$

whence

$$\tan 2\vartheta = \frac{4m_K - m_a - 3m_{f0}}{-4m_K + m_a + 3m_f}. \quad (15.14)$$

Eliminating  $\vartheta$  from Eq. (15.12) and Eq. (15.14) leads to the sum rule [4]

$$(m_f + m_{f0})(4m_K - m_a) - 3m_f m_{f0} = 8m_K^2 - 8m_K m_a + 3m_a^2. \quad (15.15)$$

**Table 15.4:**  $b$  quark-model assignments for the bottomonium and  $B$  mesons with established  $J_{PC}$ .

$n2s+1 \ J$	$J_{PC}$	$I = 0$	$I = \frac{1}{2}$	$I = 0$	$I = 0$
		$b \bar{b}$	$bu^-, bd^-;$ $\bar{b}u, \bar{b}d$	$bs^-;$ $\bar{b}s$	$bc^-;$ $\bar{b}c$
1 $1S_0$	0 <sup>+</sup>	$\eta_b(1S)$	$B$	$B_{s0}$	$B_{c\pm}$
1 $3S_1$	1 <sup>--</sup>	$\Upsilon(1S)$	$B^*$	$B_{s1}^*$	
1 $3P_0$	0 <sup>++</sup>	$\chi_{b0}(1P)$			
1 $3P_1$	1 <sup>++</sup>	$\chi_{b1}(1P)$			
1 $1P_1$	1 <sup>+-</sup>	$h_b(1P)$	$B_1(5721)$	$B_{s1}(5830)^0$	
1 $3P_2$	2 <sup>++</sup>	$\chi_{b2}(1P)$	$B_2^*(5747)$	$B_{s2}(5840)_0$	
2 $1S_0$	0 <sup>+</sup>	$\eta_b(2S)$			$B_c(2S)^\pm$

$2 \ 3S_1$	$1^{--}$	$\Upsilon(2S)$	$B_c^*(2S)_\pm$
$1 \ 3D_2$	$2^{--}$	$\Upsilon_2(1D)$	
$2 \ 3P_J$	$0, 1, 2^{++}$	$\chi_{b0,1,2}(2P)$	
$2 \ 1P_1$	$1^{+-}$	$h_b(2P)$	
$3 \ 3S_1$	$1^{--}$	$\Upsilon(3S)$	
$3 \ 3P_J$	$0, 1, 2^{++}$	$\chi_{b1,2}(3P)$	
$4 \ 3S_1$	$1^{--}$	$\Upsilon(4S)$	

This relation is verified for the ground-state vector mesons. We identify the  $\varphi(1020)$  with the  $f^0$  and the  $\omega(783)$  with the  $f$ . Thus

$$\varphi(1020) = \psi_8 \cos\vartheta_V - \psi_1 \sin\vartheta_V, \omega(783) = \psi_8 \sin\vartheta_V + \psi_1 \cos\vartheta_V, \quad (15.16)$$

with the vector mixing angle  $\vartheta_V = 36.4^\circ$  from Eq. (15.14), very close to ideal mixing. Thus  $\varphi(1020)$  is nearly pure  $ss^-$ . For ideal mixing, Eq. (15.12) and Eq. (15.14) lead to the relations

$$m_K = \frac{m_f + m_{f^0}}{2}, \quad m_\pi = m_f, \quad (15.17)$$

which are satisfied for the vector mesons.

The situation for the pseudoscalar and scalar mesons is not so clear cut, either theoretically or experimentally. For the pseudoscalars, the mixing angle is small. This can be understood qualitatively via gluon-line counting of the mixing process. The size of the mixing process between the nonstrange and strange mass bases scales as  $\alpha_s^2$ , not  $\alpha_s^3$ , because of two rather than three gluon exchange as it does for the vector mesons. It may also be that the lightest isoscalar pseudoscalars mix more strongly with excited states or with states of substantial non- $qq^-$  content, as will be discussed below.

A variety of analysis methods lead to similar results: First, for these states, Eqn. 15.15 is satisfied only approximately. Then Eqn. 15.12 and Eqn. 15.14 lead to somewhat different values for the mixing angle. Identifying the  $\eta$  with the  $f^0$  one gets

$$\eta = \psi_8 \cos\vartheta_P - \psi_1 \sin\vartheta_P, \quad (15.18)$$

$$\eta^0 = \psi_8 \sin\vartheta_P + \psi_1 \cos\vartheta_P. \quad (15.19)$$

Following chiral perturbation theory, the meson masses in the mass formulae (Eq. (15.12) and Eq. (15.14)) might be replaced by their squares. Table 15.5 lists the mixing angle  $\vartheta_{\text{lin}}$  from




Figure 15.1: SU(4) weight diagram showing the 16-plets for the pseudoscalar (a) and vector mesons (b) made of the  $u$ ,  $d$ ,  $s$ , and  $c$  quarks as a function of isospin  $I_z$ , charm  $C$ , and hypercharge  $Y$

$= B + S - \frac{C}{3}$ . The nonets of light mesons occupy the central planes to which the  $cc$  states have been added.

**Table 15.5:** Singlet-octet mixing angles for the well established nonets from the linear mass formula (15.12) and its quadratic version in which the masses are squared.

$n2s+1 J$	$JPC$	$\vartheta_{\text{quad}}$ [°]	$\vartheta_{\text{lin}}$ [°]
$11S_0$	$0^{-+}$	-11.3	-24.5
$13S_1$	$1^{--}$	39.2	36.5
$13P_2$	$2^{++}$	29.6	28.0
$13D_3$	$3^{--}$	31.8	30.8

Eqn. 15.14 (using the neutral members of the nonets) and the corresponding  $\vartheta_{\text{quad}}$  obtained by replacing the meson masses by their squares throughout.

The pseudoscalar mixing angle  $\vartheta_P$  can also be measured by comparing the partial widths for radiative  $J/\psi$  decay into a vector and a pseudoscalar [5], radiative  $\phi(1020)$  decay into  $\eta$  and  $\eta^0$  [6], radiative decays between pseudoscalar and vector mesons [7], or  $pp^-$  annihilation at rest into a pair of vector and pseudoscalar or into two pseudoscalars [8,9]. One obtains a mixing angle between  $-10^\circ$  and  $-20^\circ$ . More recently, a lattice QCD simulation, Ref. [10], has successfully reproduced the masses of the  $\eta$  and  $\eta^0$ , and as a byproduct find a mixing angle  $\vartheta_{\text{lin}} = -14.1(2.8)^\circ$ . We return to this point in Sec. 15.6.

The nonet mixing angles can be measured in  $\gamma\gamma$  collisions, *e.g.*, for the  $0^{-+}$ ,  $0^{++}$ , and  $2^{++}$  nonets. In the quark model, the amplitude for the coupling of neutral mesons to two photons is proportional to  $\sum_i Q_i^2$ , where  $Q_i$  is the charge of the  $i$ -th quark. The  $2\gamma$  partial width of an isoscalar meson with mass  $m$  is then given in terms of the mixing angle  $\alpha$  by

$$\Gamma_{2\gamma} = C(5\cos\alpha - \sqrt{2}\sin\alpha)^2 m^3, \quad (15.20)$$

- for  $f^0$  and  $f$  ( $\alpha \rightarrow \alpha - 90^\circ$ ). The coupling  $C$  may depend on the meson mass. It is often assumed to be a constant in the nonet. For the isovector  $a$ , one then finds  $\Gamma_{2\gamma} = 9 C m^3$ . Thus the members of an ideally mixed nonet couple to  $2\gamma$  with partial widths in the ratios  $f : f^0 : a = 25 : 2 : 9$ . For tensor mesons, one finds from the ratios of the measured  $2\gamma$  partial widths for the  $f_2(1270)$  and  $f_2^0(1525)$  mesons a mixing angle  $\alpha_T$  of  $(81 \pm 1)^\circ$ , or  $\vartheta_T = (27 \pm 1)^\circ$ , in accord with the linear mass formula. For the pseudoscalars, one finds from the ratios of partial widths  $\Gamma(\eta^0 \rightarrow 2\gamma)/\Gamma(\eta \rightarrow 2\gamma)$  a mixing angle  $\vartheta_P = (-18 \pm 2)^\circ$ , while the ratio  $\Gamma(\eta^0 \rightarrow 2\gamma)/\Gamma(\pi^0 \rightarrow 2\gamma)$  leads to  $\sim -24^\circ$ . SU(3) breaking effects for pseudoscalars are discussed in [11].

The partial width for the decay of a scalar or a tensor meson into a pair of pseudoscalar mesons is model-dependent. Following Ref. [12],

$$\Gamma = C \times \gamma^2 \times |F(q)|^2 \times q. \quad (15.21)$$

$C$  is a nonet constant,  $q$  the momentum of the decay products,  $F(q)$  a form factor, and  $\gamma^2$  the SU(3) coupling. The model-dependent form factor may be written as

$$|F(q)|^2 = q^2 \times \exp\left(-\frac{q^2}{8\theta^2}\right), \quad (15.22)$$

where  $\ell$  is the relative angular momentum between the decay products. The decay of a  $q\bar{q}$  meson into a pair of mesons involves the creation of a  $q\bar{q}$  pair from the vacuum, and SU(3) symmetry assumes that the matrix elements for the creation of  $s\bar{s}$ ,  $u\bar{u}$ , and  $d\bar{d}$  pairs are equal. The couplings  $\gamma^2$  are given in Table 15.6, and their dependence upon the mixing angle  $\alpha$  is shown in Fig. 15.2 for isoscalar decays. The generalization to unequal  $s\bar{s}$ ,  $u\bar{u}$ , and  $d\bar{d}$  couplings is given in Ref. [12]. An excellent fit to the tensor meson decay widths is obtained assuming SU(3) symmetry, with  $\theta \approx 0.5 \text{ GeV/c}$ ,  $\vartheta_V \approx 26^\circ$  and  $\vartheta_P \approx -17^\circ$  [12].

**Table 15.6:** SU(3) couplings  $\gamma^2$  for quarkonium decays as a function of nonet mixing angle  $\alpha$ , up to a common multiplicative factor  $C$  ( $\varphi = 54.7^\circ + \vartheta_P$ ).

Isospin	Decay channel	$\gamma^2$
0	$\pi\pi$	$3 \cos^2 \alpha$
	$K\bar{K}$	$(\cos \alpha - \frac{1}{\sqrt{2}} \sin \alpha)^2$
	$\eta\eta$	$(\cos \alpha \cos^2 \varphi - \frac{1}{2} \sin^2 2\varphi \cos \alpha + \frac{1}{2} \sin^2 \alpha \sin^2 \varphi)^2$
	$\eta\eta^0$	$\frac{1}{2} \sin^2 2\varphi (\cos \alpha + \frac{1}{\sqrt{2}} \sin \alpha)^2$
1	$\eta\pi$	$2 \cos^2 \varphi$
	$\eta^0\pi$	$2 \sin^2 \varphi$
	$K\bar{K}$	1
$\frac{1}{2}$	$K\pi$	$\frac{3}{2} (\sin \varphi - \frac{\cos \varphi}{\sqrt{2}})^2$
	$K\eta$	$(\sin \varphi - \frac{\cos \varphi}{\sqrt{2}})^2$
	$K\eta^0$	$(\cos \varphi + \frac{\sin \varphi}{\sqrt{2}})^2$




Figure 15.2: SU(3) couplings as a function of mixing angle  $\alpha$  for isoscalar decays, up to a common multiplicative factor  $C$  and for  $\vartheta_p = -17.3^\circ$ .

### 15.3 Exotic mesons

The existence of a light nonet composed of four quarks (tetraquarks) with masses below 1 GeV was suggested a long time ago [13] [14]. Coupling two triplets of light quarks  $u$ ,  $d$ , and  $s$ , one obtains nine states, of which the six symmetric ( $uu$ ,  $dd$ ,  $ss$ ,  $ud + du$ ,  $us + su$ ,  $ds + sd$ ) form the six dimensional representation **6**, while the three antisymmetric ( $ud - du$ ,  $us - su$ ,  $ds - sd$ ) form

the three dimensional representation **3** of SU(3):

$$\mathbf{3} \otimes \mathbf{3} = \mathbf{6} \oplus \bar{\mathbf{3}}. \quad (15.23)$$

Hence for tetraquarks one gets the reduction

$$\begin{aligned} & \mathbf{3} \otimes \mathbf{3} \otimes \bar{\mathbf{3}} \otimes \bar{\mathbf{3}} \\ &= \mathbf{6} \oplus \bar{\mathbf{3}} \otimes \bar{\mathbf{6}} \otimes \mathbf{3} \\ &= \bar{\mathbf{3}} \otimes \mathbf{3} \oplus \mathbf{6} \otimes \bar{\mathbf{6}} \oplus \mathbf{6} \otimes \mathbf{3} \oplus \bar{\mathbf{3}} \otimes \bar{\mathbf{6}} \\ &= \mathbf{9} \oplus \mathbf{36} \oplus \mathbf{18} \oplus \bar{\mathbf{18}}. \end{aligned} \quad (15.24)$$


$$(15.25)$$

Combining with spin and color and requiring antisymmetry for diquarks and antidiquarks, one finds for ground states (zero angular momenta) that the most deeply bound tetraquarks (and hence the lightest ones) lie in the nonet and are scalar mesons (see also [1]). The average mass is estimated to be around 900 MeV from the mass differences between the  $\rho$  and  $\pi$  masses. Letting the strange quark determine the mass splittings one obtains a mass inverted spectrum with a light isosinglet, a medium heavy isodoublet and a heavy isotriplet + isosinglet. It is then tempting to identify these mesons as the lightest scalars

$$\begin{aligned} f_0(500) &= \bar{u}dud, K^-(700) = (\bar{s}dud, \bar{s}ud) \text{ and } (\bar{u}dus, \bar{u}dds), \\ a_0(980) &= (us\bar{d}\bar{s}, \frac{1}{\sqrt{2}}[u\bar{u} - d\bar{d}]s\bar{s}, \bar{u}\bar{s}ds), f_0(980) = \frac{1}{\sqrt{2}}[u\bar{u} + d\bar{d}]s\bar{s}. \end{aligned} \quad (15.26)$$

A plethora of new states have been reported in the charmonium and bottomonium spectra. The most prominent one is the  $\chi_{c1}(3872)$  (formerly  $X(3872)$ ), first observed in 2003 in  $B$ -decays in the final state  $J/\psi \pi^+ \pi^-$  (see Fig. 15.3). Even more remarkable is the observation of isovector (charged) mesons decaying into  $cc$  plus a charged pion, such as the  $Z^+(4430)$  decaying into  $\psi(2S)\pi^+$ , which a priori excludes an interpretation as true  $cc$  (charmonium) state. Similar states are also observed in the bottomonium spectrum. Some of these states may be tetraquarks (e.g.  $cqc^-q^-$ ), molecular structures (e.g.  $cq^-cq^-$ ) made of pairs of mesons such as  $D$ ,  $D_s$  and  $D^*$ ,  $D_s^*$  excitations, or their  $B$  and  $B^*$  counterparts. They could also be mimicked by kinematical effects. Details and references can be found in recent reviews [15], [16] and in the “Note on Non- $qq^-$  Mesons” in the Meson Listings.

QCD predicts the existence of extra isoscalar mesons. In the pure gauge theory they contain only gluons, and are called glueballs. The ground state glueball is predicted by lattice gauge theories to be  $0^{++}$ , the first excited state  $2^{++}$ . Errors on the mass predictions are large. From Ref. [17] one obtains 1750 (50) (80) MeV for the mass of the lightest  $0^{++}$  glueball from quenched QCD. As an example for the glueball mass



spectrum, we show in Fig. 15.4 a calculation from Ref. [18]. A mass of 1710 MeV is predicted for the ground state, also with an error of about 100 MeV. Earlier work by other groups produced masses at 1650 MeV [19] and 1550 MeV [20] (see also [21]). The first excited state has a mass of about 2.4 GeV, and the lightest glueball with exotic quantum numbers ( $2^{+-}$ ) has a mass of about 4 GeV.

State	Mass (MeV)	Width (MeV)	Decay
$\psi(3770)$	3770	11	$D_s^+ D_s^-$
$\psi(4040)$	4040	10	$D_s^+ D_s^-$
$\psi(4180)$	4180	10	$D_s^+ D_s^-$
$\psi(4415)$	4415	10	$D_s^+ D_s^-$
$\psi(4730)$	4730	10	$D_s^+ D_s^-$
$\psi(5040)$	5040	10	$D_s^+ D_s^-$
$\psi(5400)$	5400	10	$D_s^+ D_s^-$
$\psi(5720)$	5720	10	$D_s^+ D_s^-$
$\psi(6040)$	6040	10	$D_s^+ D_s^-$
$\psi(6350)$	6350	10	$D_s^+ D_s^-$
$\psi(6670)$	6670	10	$D_s^+ D_s^-$
$\psi(6980)$	6980	10	$D_s^+ D_s^-$
$\psi(7290)$	7290	10	$D_s^+ D_s^-$
$\psi(7610)$	7610	10	$D_s^+ D_s^-$
$\psi(7930)$	7930	10	$D_s^+ D_s^-$
$\psi(8250)$	8250	10	$D_s^+ D_s^-$
$\psi(8570)$	8570	10	$D_s^+ D_s^-$
$\psi(8890)$	8890	10	$D_s^+ D_s^-$
$\psi(9210)$	9210	10	$D_s^+ D_s^-$
$\psi(9530)$	9530	10	$D_s^+ D_s^-$
$\psi(9850)$	9850	10	$D_s^+ D_s^-$
$\psi(10170)$	10170	10	$D_s^+ D_s^-$
$\psi(10490)$	10490	10	$D_s^+ D_s^-$
$\psi(10810)$	10810	10	$D_s^+ D_s^-$
$\psi(11130)$	11130	10	$D_s^+ D_s^-$
$\psi(11450)$	11450	10	$D_s^+ D_s^-$
$\psi(11770)$	11770	10	$D_s^+ D_s^-$
$\psi(12090)$	12090	10	$D_s^+ D_s^-$
$\psi(12410)$	12410	10	$D_s^+ D_s^-$
$\psi(12730)$	12730	10	$D_s^+ D_s^-$
$\psi(13050)$	13050	10	$D_s^+ D_s^-$
$\psi(13370)$	13370	10	$D_s^+ D_s^-$
$\psi(13690)$	13690	10	$D_s^+ D_s^-$
$\psi(14010)$	14010	10	$D_s^+ D_s^-$
$\psi(14330)$	14330	10	$D_s^+ D_s^-$
$\psi(14650)$	14650	10	$D_s^+ D_s^-$
$\psi(14970)$	14970	10	$D_s^+ D_s^-$
$\psi(15290)$	15290	10	$D_s^+ D_s^-$
$\psi(15610)$	15610	10	$D_s^+ D_s^-$
$\psi(15930)$	15930	10	$D_s^+ D_s^-$
$\psi(16250)$	16250	10	$D_s^+ D_s^-$
$\psi(16570)$	16570	10	$D_s^+ D_s^-$
$\psi(16890)$	16890	10	$D_s^+ D_s^-$
$\psi(17210)$	17210	10	$D_s^+ D_s^-$
$\psi(17530)$	17530	10	$D_s^+ D_s^-$
$\psi(17850)$	17850	10	$D_s^+ D_s^-$
$\psi(18170)$	18170	10	$D_s^+ D_s^-$
$\psi(18490)$	18490	10	$D_s^+ D_s^-$
$\psi(18810)$	18810	10	$D_s^+ D_s^-$
$\psi(19130)$	19130	10	$D_s^+ D_s^-$
$\psi(19450)$	19450	10	$D_s^+ D_s^-$
$\psi(19770)$	19770	10	$D_s^+ D_s^-$
$\psi(20090)$	20090	10	$D_s^+ D_s^-$
$\psi(20410)$	20410	10	$D_s^+ D_s^-$
$\psi(20730)$	20730	10	$D_s^+ D_s^-$
$\psi(21050)$	21050	10	$D_s^+ D_s^-$
$\psi(21370)$	21370	10	$D_s^+ D_s^-$
$\psi(21690)$	21690	10	$D_s^+ D_s^-$
$\psi(22010)$	22010	10	$D_s^+ D_s^-$
$\psi(22330)$	22330	10	$D_s^+ D_s^-$
$\psi(22650)$	22650	10	$D_s^+ D_s^-$
$\psi(22970)$	22970	10	$D_s^+ D_s^-$
$\psi(23290)$	23290	10	$D_s^+ D_s^-$
$\psi(23610)$	23610	10	$D_s^+ D_s^-$
$\psi(23930)$	23930	10	$D_s^+ D_s^-$
$\psi(24250)$	24250	10	$D_s^+ D_s^-$
$\psi(24570)$	24570	10	$D_s^+ D_s^-$
$\psi(24890)$	24890	10	$D_s^+ D_s^-$
$\psi(25210)$	25210	10	$D_s^+ D_s^-$
$\psi(25530)$	25530	10	$D_s^+ D_s^-$
$\psi(25850)$	25850	10	$D_s^+ D_s^-$
$\psi(26170)$	26170	10	$D_s^+ D_s^-$
$\psi(26490)$	26490	10	$D_s^+ D_s^-$
$\psi(26810)$	26810	10	$D_s^+ D_s^-$
$\psi(27130)$	27130	10	$D_s^+ D_s^-$
$\psi(27450)$	27450	10	$D_s^+ D_s^-$
$\psi(27770)$	27770	10	$D_s^+ D_s^-$
$\psi(28090)$	28090	10	$D_s^+ D_s^-$
$\psi(28410)$	28410	10	$D_s^+ D_s^-$
$\psi(28730)$	28730	10	$D_s^+ D_s^-$
$\psi(29050)$	29050	10	$D_s^+ D_s^-$
$\psi(29370)$	29370	10	$D_s^+ D_s^-$
$\psi(29690)$	29690	10	$D_s^+ D_s^-$
$\psi(30010)$	30010	10	$D_s^+ D_s^-$
$\psi(30330)$	30330	10	$D_s^+ D_s^-$
$\psi(30650)$	30650	10	$D_s^+ D_s^-$
$\psi(30970)$	30970	10	$D_s^+ D_s^-$
$\psi(31290)$	31290	10	$D_s^+ D_s^-$
$\psi(31610)$	31610	10	$D_s^+ D_s^-$
$\psi(31930)$	31930	10	$D_s^+ D_s^-$
$\psi(32250)$	32250	10	$D_s^+ D_s^-$
$\psi(32570)$	32570	10	$D_s^+ D_s^-$
$\psi(32890)$	32890	10	$D_s^+ D_s^-$
$\psi(33210)$	33210	10	$D_s^+ D_s^-$
$\psi(33530)$	33530	10	$D_s^+ D_s^-$
$\psi(33850)$	33850	10	$D_s^+ D_s^-$
$\psi(34170)$	34170	10	$D_s^+ D_s^-$
$\psi(34490)$	34490	10	$D_s^+ D_s^-$
$\psi(34810)$	34810	10	$D_s^+ D_s^-$
$\psi(35130)$	35130	10	$D_s^+ D_s^-$
$\psi(35450)$	35450	10	$D_s^+ D_s^-$
$\psi(35770)$	35770	10	$D_s^+ D_s^-$
$\psi(36090)$	36090	10	$D_s^+ D_s^-$
$\psi(36410)$	36410	10	$D_s^+ D_s^-$
$\psi(36730)$	36730	10	$D_s^+ D_s^-$
$\psi(37050)$	37050	10	$D_s^+ D_s^-$
$\psi(37370)$	37370	10	$D_s^+ D_s^-$
$\psi(37690)$	37690	10	$D_s^+ D_s^-$
$\psi(38010)$	38010	10	$D_s^+ D_s^-$
$\psi(38330)$	38330	10	$D_s^+ D_s^-$
$\psi(38650)$	38650	10	$D_s^+ D_s^-$
$\psi(38970)$	38970	10	$D_s^+ D_s^-$
$\psi(39290)$	39290	10	$D_s^+ D_s^-$
$\psi(39610)$	39610	10	$D_s^+ D_s^-$
$\psi(39930)$	39930	10	$D_s^+ D_s^-$
$\psi(40250)$	40250	10	$D_s^+ D_s^-$
$\psi(40570)$	40570	10	$D_s^+ D_s^-$
$\psi(40890)$	40890	10	$D_s^+ D_s^-$
$\psi(41210)$	41210	10	$D_s^+ D_s^-$
$\psi(41530)$	41530	10	$D_s^+ D_s^-$
$\psi(41850)$	41850	10	$D_s^+ D_s^-$
$\psi(42170)$	42170	10	$D_s^+ D_s^-$
$\psi(42490)$	42490	10	$D_s^+ D_s^-$
$\psi(42810)$	42810	10	$D_s^+ D_s^-$
$\psi(43130)$	43130	10	$D_s^+ D_s^-$
$\psi(43450)$	43450	10	$D_s^+ D_s^-$
$\psi(43770)$	43770	10	$D_s^+ D_s^-$
$\psi(44090)$	44090	10	$D_s^+ D_s^-$
$\psi(44410)$	44410	10	$D_s^+ D_s^-$
$\psi(44730)$	44730	10	$D_s^+ D_s^-$
$\psi(45050)$	45050	10	$D_s^+ D_s^-$
$\psi(45370)$	45370	10	$D_s^+ D_s^-$
$\psi(45690)$	45690	10	$D_s^+ D_s^-$
$\psi(46010)$	46010	10	$D_s^+ D_s^-$
$\psi(46330)$	46330	10	$D_s^+ D_s^-$
$\psi(46650)$	46650	10	$D_s^+ D_s^-$
$\psi(46970)$	46970	10	$D_s^+ D_s^-$
$\psi(47290)$	47290	10	$D_s^+ D_s^-$
$\psi(47610)$	47610	10	$D_s^+ D_s^-$
$\psi(47930)$	47930	10	$D_s^+ D_s^-$
$\psi(48250)$	48250	10	$D_s^+ D_s^-$
$\psi(48570)$	48570	10	$D_s^+ D_s^-$
$\psi(48890)$	48890	10	$D_s^+ D_s^-$
$\psi(49210)$	49210	10	$D_s^+ D_s^-$
$\psi(49530)$	49530	10	$D_s^+ D_s^-$
$\psi(49850)$	49850	10	$D_s^+ D_s^-$
$\psi(50170)$	50170	10	$D_s^+ D_s^-$
$\psi(50490)$	50490	10	$D_s^+ D_s^-$
$\psi(50810)$	50810	10	$D_s^+ D_s^-$
$\psi(51130)$	51130	10	$D_s^+ D_s^-$
$\psi(51450)$	51450	10	$D_s^+ D_s^-$
$\psi(51770)$	51770	10	$D_s^+ D_s^-$
$\psi(52090)$	52090	10	$D_s^+ D_s^-$
$\psi(52410)$	52410	10	$D_s^+ D_s^-$
$\psi(52730)$	52730	10	$D_s^+ D_s^-$
$\psi(53050)$	53050	10	$D_s^+ D_s^-$
$\psi(53370)$	53370	10	$D_s^+ D_s^-$
$\psi(53690)$	53690	10	$D_s^+ D_s^-$
$\psi(54010)$	54010	10	$D_s^+ D_s^-$
$\psi(54330)$	54330	10	$D_s^+ D_s^-$
$\psi(54650)$	54650	10	$D_s^+ D_s^-$
$\psi(54970)$	54970	10	$D_s^+ D_s^-$
$\psi(55290)$	55290	10	$D_s^+ D_s^-$
$\psi(55610)$	55610	10	$D_s^+ D_s^-$
$\psi(55930)$	55930	10	$D_s^+ D_s^-$
$\psi(56250)$	56250	10	$D_s^+ D_s^-$
$\psi(56570)$	56570	10	$D_s^+ D_s^-$
$\psi(56890)$	56890	10	$D_s^+ D_s^-$
$\psi(57210)$	57210	10	$D_s^+ D_s^-$
$\psi(57530)$	57530	10	$D_s^+ D_s^-$
$\psi(57850)$	57850	10	$D_s^+ D_s^-$
$\psi(58170)$	58170	10	$D_s^+ D_s^-$
$\psi(58490)$	58490	10	$D_s^+ D_s^-$
$\psi(58810)$	58810	10	$D_s^+ D_s^-$
$\psi(59130)$	59130	10	$D_s^+ D_s^-$
$\psi(59450)$	59450	10	$D_s^+ D_s^-$
$\psi(59770)$	59770	10	$D_s^+ D_s^-$
$\psi(60090)$	60090	10	$D_s^+ D_s^-$
$\psi(60410)$	60410	10	$D_s^+ D_s^-$
$\psi(60730)$	60730	10	$D_s^+ D_s^-$
$\psi(61050)$	61050	10	$D_s^+ D_s^-$
$\psi(61370)$	61370	10	$D_s^+ D_s^-$
$\psi(61690)$	61690	10	$D_s^+ D_s^-$
$\psi(62010)$	62010	10	$D_s^+ D_s^-$
$\psi(62330)$	62330	10	$D_s^+ D_s^-$
$\psi(62650)$	62650	10	$D_s^+ D_s^-$
$\psi(62970)$	62970	10	$D_s^+ D_s^-$
$\psi(63290)$	63290	10	$D_s^+ D_s^-$
$\psi(63610)$	63610	10	$D_s^+ D_s^-$
$\psi(63930)$	63930	10	$D_s^+ D_s^-$
$\psi(64250)$	64250	10	$D_s^+ D_s^-$
$\psi(64570)$	64570	10	$D_s^+ D_s^-$
$\psi(64890)$	64890	10	$D_s^+ D_s^-$
$\psi(65210)$	65210	10	$D_s^+ D_s^-$
$\psi(65530)$	65530	10	$D_s^+ D_s^-$
$\psi(65850)$	65850	10	$D_s^+ D_s^-$
$\psi(66170)$	66170	10	$D_s^+ D_s^-$
$\psi(66490)$	66490	10	$D_s^+ D_s^-$
$\psi(66810)$	66810	10	$D_s^+ D_s^-$
$\psi(67130)$	67130	10	$D_s^+ D_s^-$
$\psi(67450)$	67450	10	$D_s^+ D_s^-$
$\psi(67770)$	67770	10	$D_s^+ D_s^-$
$\psi(68090)$	68090	10	$D_s^+ D_s^-$
$\psi(68410)$	68410	10	$D_s^+ D_s^-$
$\psi(68730)$	68730	10	$D_s^+ D_s^-$
$\psi(69050)$	69050	10	$D_s^+ D_s^-$
$\psi(69370)$	69370	10	$D_s^+ D_s^-$
$\psi(69690)$	69690	10	$D_s^+ D_s^-$
$\psi(69990)$	69990	10	$D_s^+ D_s^-$
$\psi(70290)$	70290	10	$D_s^+ D_s^-$
$\psi(70590)$	70590	10	$D_s^+ D_s^-$
$\psi(70890)$	70890	10	$D_s^+ D_s^-$
$\psi(71190)$	71190	10	$D_s^+ D_s^-$
$\psi(71490)$	71490	10	$D_s^+ D_s^-$
$\psi(71790)$	71790	10	$D_s^+ D_s^-$
$\psi(72090)$	72090	10	$D_s^+ D_s^-$
$\psi(72390)$	72390	10	$D_s^+ D_s^-$
$\psi(72690)$	72690	10	$D_s^+ D_s^-$
$\psi(72990)$	72990	10	$D_s^+ D_s^-$
$\psi(73290)$	73290	10	$D_s^+ D_s^-$
$\psi(73590)$	73590	10	$D_s^+ D_s^-$
$\psi(73890)$	73890	10	$D_s^+ D_s^-$
$\psi(74190)$	74190	10	$D_s^+ D_s^-$
$\psi(74490)$	74490	10	$D_s^+ D_s^-$
$\psi(74790)$	74790	10	$D_s^+ D_s^-$
$\psi(75090)$	75090	10	$D_s^+ D_s^-$
$\psi(75390)$	75390	10	$D_s^+ D_s^-$
$\psi(75690)$	75690	10	




Figure 15.4: Predicted glueball mass spectrum from the lattice in quenched approximation (from [18]).

These calculations are made in the so-called “quenched approximation” which neglects  $qq^-$  loops. However, both glue and  $qq^-$  states couple to singlet scalar mesons. Therefore glueballs will mix with nearby  $qq^-$  states of the same quantum numbers. For example, the two isoscalar  $0^{++}$  mesons around 1500 MeV will mix with the pure ground state glueball to generate the observed physical states  $f_0(1370)$ ,  $f_0(1500)$ , and  $f_0(1710)$  [12,22]. The first results from lattice calculations, which include these effects, indicate that the mass shifts are small. We return to a discussion of this point in Sec. 15.6.

The existence of three singlet scalar mesons around 1.5 GeV suggests additional degrees of freedom such as glue, since only two mesons are predicted in this mass range. The  $f_0(1500)$  [12,22] or, alternatively, the  $f_0(1710)$  [19], have been proposed as candidates for the scalar glueball, both states having considerable mixing also with the  $f_0(1370)$ . Other mixing schemes, in particular with the  $f_0(500)$  and the  $f_0(980)$ , have also been proposed [23]. According to a holographic model of low-energy QCD scalar glueballs decay strongly into kaons and  $\eta$  mesons, in good agreement with data on the  $f_0(1710)$  [24]. Details can be found in the “Note on Non- $qq^-$  Mesons” in the Meson Listings and in Ref. [25]. See also the “Note on Scalar Mesons below 2 GeV”.

Mesons made of  $qq^-$  pairs bound by excited gluons  $g$ , the hybrid states  $qqg^-$ , are also predicted. They should lie in the 1.9 GeV mass region, according to gluon flux tube models [26]. Lattice QCD also predicts

the lightest hybrid, an exotic  $1^{-+}$ , at a mass of 1.8 to 1.9 GeV [27]. However, the bag model predicts four nonets, among them an exotic  $1^{-+}$  around or above 1.4 GeV [28,29]. There are so far two candidates for exotic states with quantum numbers  $1^{-+}$ , the  $\pi_1(1400)$  and  $\pi_1(1600)$ , which could be hybrids or four-quark states (see the “Note on Non- $qq^{-}$  Mesons” in the Meson Listings and in [25]).

#### 15.4 Baryons: $qqq$ states

Baryons are fermions with baryon number  $B = 1$ , *i.e.*, in the most general case, they are composed of three quarks plus any number of quark - antiquark pairs. Until recently, all established baryons were 3-quark ( $qqq$ ) configurations, which we mainly discuss in this section. However, in 2015 the LHCb collaboration published first evidence for charmed ‘pentaquark’ states of minimal quark content  $ccuud^{-}$  at invariant masses close to 4.4 GeV [30.] More refined LHCb experiments have revealed evidence for three such states called  $P_c(4312)^+$ ,  $P_c(4440)^+$ , and  $P_c(4457)^+$  [31]. These states are located close to the thresholds of the production of ordinary baryon-meson pairs like  $\Sigma_c^+ \bar{D}^0$  and  $\Sigma_c^+ \bar{D}^{*0}$  and are discussed in terms of molecular-like states. A nice overview on the discussion of pentaquark and tetraquark states is given in Ref. [32].

The color part of baryon state functions is an SU(3) singlet, a completely antisymmetric state of the three colors. Since the quarks are fermions, the state function must be antisymmetric under interchange of any two equal-mass quarks (up and down quarks in the limit of isospin symmetry). Thus it can be written as

$$|qqq\rangle_{i_A} = |\text{color}\rangle_{i_A} \times |\text{space, spin, flavor}\rangle_s, \quad (15.27)$$

where the subscripts  $S$  and  $A$  indicate symmetry or antisymmetry under interchange of any two equal-mass quarks. Note the contrast with the state function for the three nucleons in  $^3\text{H}$  or  $^3\text{He}$ :

$$|NNN\rangle_{i_A} = |\text{space, spin, isospin}\rangle_{i_A}. \quad (15.28)$$

This difference has major implications for internal structure, magnetic moments, *etc.* (For a nice discussion, see Ref. [33].)

The “ordinary” baryons are made up of  $u$ ,  $d$ , and  $s$  quarks. The three flavors imply an approximate flavor SU(3), which requires that baryons made of these quarks belong to the multiplets on the right side of

$$\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{10}_S \oplus \mathbf{8}_M \oplus \mathbf{8}_M \oplus \mathbf{1}_A \quad (15.29)$$

(see the section on “SU( $n$ ) Multiplets and Young Diagrams”). Here the subscripts indicate symmetric, mixed-symmetry, or antisymmetric states under interchange of any two quarks. The  $\mathbf{1}$  is a  $uds$  state ( $\Lambda_1$ ), and the octet contains a similar state ( $\Lambda_8$ ). If these have the same spin and parity, they can mix. The mechanism is the same as for the mesons (see above). In the ground state multiplet, the SU(3) flavor singlet  $\Lambda_1$  is forbidden by Fermi statistics. The section on “SU(3) Isoscalar Factors and Representation Matrices,” shows how relative decay rates in, say,  $\mathbf{10} \rightarrow \mathbf{8} \otimes \mathbf{8}$  decays may be calculated.

The addition of the  $c$  quark to the light quarks extends the flavor symmetry to SU(4). However, due to the large mass of the  $c$  quark, this symmetry is much more strongly broken than the SU(3) of the three light quarks. Figures 15.5(a) and 15.5(b) show the SU(4) baryon multiplets that have as their bottom levels an SU(3) octet, such as the octet that includes the nucleon, or an SU(3) decuplet, such as the decuplet that includes the  $\Delta(1232)$ . All particles in a given SU(4) multiplet have the same spin and parity. The charmed baryons are discussed in more detail in the “Note on Charmed Baryons” in the Particle Listings. The same

multiplets as shown in Fig. 15.5 can be constructed when the  $c$  quark is replaced by the  $b$  quark, or they can be embedded in a larger SU(5) group that accounts for all baryons that can be constructed from the five quark flavors. The existence of baryons with  $t$ -quarks is very unlikely due to the short lifetime of the  $t$ -quark. The heavy quark baryons have recently gained a lot of interest [34]. Their relatively narrow widths allow to isolate the states much easier than the light quark baryon resonances which require intricate partial wave analyses. The only problem on the experimental side are the small production cross sections, but the recent measurements at the  $e^+e^-$  colliding  $B$  factories, at the  $pp^-$  Tevatron collider, and at LHCb at CERN have boosted this field. The LHCb collaboration has published evidence for five new narrow  $\Omega_c^0$  states ( $css$ ) [35] and for a doubly charmed  $\Xi_{cc}^{++}$  ( $ccu$ ) [36] baryon. Doubly charmed baryons have a much different structure from light baryons, more resembling a heavy ‘double-star’ system with an attached light ‘planet’ and open a new window for QCD properties. Another candidate for a doubly charmed baryon ( $\Xi_{cc}^+$ , ( $ccd$ )) had been earlier reported by the SELEX experiment [37,38] but could so far not be confirmed by other experiments and the difference in mass between the LHCb  $\Xi_{cc}^{++}$  and the SELEX  $\Xi_{cc}^+$  would be much larger than predicted. Quark model predictions for baryons with two heavy quarks are given in Ref. [39] and lattice results for doubly and triply charmed states are discussed in Sec. 15.6 of this review.

For the “ordinary” baryons (no  $c$  or  $b$  quark), flavor and spin may be combined in an approximate flavor-spin SU(6), in which the six basic states are  $d \uparrow$ ,  $d \downarrow$ , ...,  $s \downarrow$  ( $\uparrow$ ,  $\downarrow$  = spin up, down). Then the baryons belong to the multiplets on the right side of

$$6 \otimes 6 \otimes 6 = 56_S \oplus 70_M \oplus 70_M \oplus 20_A.$$

These SU(6) multiplets decompose into flavor SU(3) multiplets as follows:

$$56 = {}^4 10 \oplus {}^2 8 \quad (15.30)$$

$$70 = {}^2 10 \oplus {}^4 8 \oplus {}^2 8 \oplus {}^2 1 \quad (15.31)$$

$$20 = {}^2 8 \oplus {}^4 1, \quad (15.32)$$

where the superscript ( $2S + 1$ ) gives the net spin  $S$  of the quarks for each particle in the SU(3) multiplet. The  $J^P = 1/2^+$  octet containing the nucleon and the  $J^P = 3/2^+$  decuplet containing the  $\Delta(1232)$  together make up the “ground-state” 56-plet, in which the orbital angular momenta between the quark pairs are zero (so that the spatial part of the state function is trivially symmetric). The **70** and **20** require some excitation of the spatial part of the state function in order to make the overall state function symmetric. States with nonzero orbital angular momenta are classified in SU(6)  $\otimes$  O(3) supermultiplets.

It is useful to classify the baryons into bands that have the same number  $N$  of quanta of excitation. Each band consists of a number of supermultiplets, specified by  $(D, L^P_N)$ , where  $D$  is the dimensionality of the SU(6) representation,  $L$  is the total quark orbital angular momentum, and  $P$  is the total parity. Supermultiplets contained in bands up to  $N = 12$  are given in Ref. [40]. The  $N = 0$  band, which contains the nucleon and  $\Delta(1232)$ , consists only of the  $(56, 0^+_0)$  supermultiplet. The  $N = 1$  band consists only of the  $(70, 1^-_1)$  multiplet and contains the negative-parity baryons with masses below about 1.9 GeV. The  $N = 2$  band contains five supermultiplets:  $(56, 0^+_2)$ ,  $(70, 0^+_2)$ ,  $(56, 2^+_2)$ ,  $(70, 2^+_2)$ , and  $(20, 1^+_2)$ .

The wave functions of the non-strange baryons in the harmonic oscillator basis are often labeled by  $|X^{2S+1} L_\pi J^P i\rangle$ , where  $S, L, J, P$  are as above,  $X = N$  or  $\Delta$ , and  $\pi = S, M$  or  $A$  denotes the symmetry of the spatial

wave function. The possible model states for the bands with  $N=0,1,2$  are given in Table 15.8. The assignment of experimentally observed states is only complete and




Figure 15.5:  $SU(4)$  multiplets of baryons made of  $u$ ,  $d$ ,  $s$ , and  $c$  quarks. (a) The 20-plet with an  $SU(3)$  octet. (b) The 20-plet with an  $SU(3)$  decuplet.

well established up to the  $N=1$  band. Some more tentative assignments for higher multiplets are suggested in [41].

In Table 15.7, quark-model assignments are given for many of the established baryons whose  $SU(6) \otimes O(3)$  compositions are relatively unmixed. One must, however, keep in mind that apart from the

mixing of the  $\Lambda$  singlet and octet states, states with same  $J^P$  but different  $L, S$  combinations can also mix. In the quark model with one-gluon exchange motivated interactions, the size of the mixing is determined by the relative strength of the tensor term with respect to the contact term **Table 15.7:** Quark-model assignments for some of the known baryons in terms of a flavor-spin SU(6) basis. Only the dominant representation is listed. Assignments for several states, especially for the  $\Lambda(1810)$ ,  $\Lambda(2350)$ ,  $\Xi(1820)$ , and  $\Xi(2030)$ , are merely educated guesses.

<sup>†</sup>suggestions for assignments and re-assignments from Ref. [42]. For assignments of the charmed baryons, see the “Note on Charmed Baryons” in the Particle Listings.

$J^P$	$(D, L^P_N)$	$S$	Octet members				Singlets
$1/2^+$	$(56, 0^+_0)$	$1/2$	$N(939)$	$\Lambda(1116)$	$\Sigma(1193)$	$\Xi(1318)$	
$1/2^+$	$(56, 0^+_2)$	$1/2$	$N(1440)$	$\Lambda(1600)$	$\Sigma(1660)$	$\Xi(1690)^{\dagger}$	
$1/2^-$	$(70, 1^-_1)$	$1/2$	$N(1535)$	$\Lambda(1670)$	$\Sigma(1620)$ $\Sigma(1560)^{\dagger}$	$\Xi(?)$	$\Lambda(1405)$
$3/2^-$	$(70, 1^-_1)$	$1/2$	$N(1520)$	$\Lambda(1690)$	$\Sigma(1670)$	$\Xi(1820)$	$\Lambda(1520)$
$1/2^-$	$(70, 1^-_1)$	$3/2$	$N(1650)$	$\Lambda(1800)$	$\Sigma(1750)$ $\Sigma(1620)^{\dagger}$	$\Xi(?)$	
$3/2^-$	$(70, 1^-_1)$	$3/2$	$N(1700)$	$\Lambda(?)$	$\Sigma(1940)^{\dagger}$	$\Xi(?)$	
$5/2^-$	$(70, 1^-_1)$	$3/2$	$N(1675)$	$\Lambda(1830)$	$\Sigma(1775)$	$\Xi(1950)^{\dagger}$	
$1/2^+$	$(70, 0^+_2)$	$1/2$	$N(1710)$	$\Lambda(1810)$	$\Sigma(1880)$	$\Xi(?)$	$\Lambda(1810)^{\dagger}$
$3/2^+$	$(56, 2^+_2)$	$1/2$	$N(1720)$	$\Lambda(1890)$	$\Sigma(?)$	$\Xi(?)$	
$5/2^+$	$(56, 2^+_2)$	$1/2$	$N(1680)$	$\Lambda(1820)$	$\Sigma(1915)$	$\Xi(2030)$	
$7/2^-$	$(70, 3^-_3)$	$1/2$	$N(2190)$	$\Lambda(?)$	$\Sigma(?)$	$\Xi(?)$	$\Lambda(2100)$
$9/2^-$	$(70, 3^-_3)$	$3/2$	$N(2250)$	$\Lambda(?)$	$\Sigma(?)$	$\Xi(?)$	
$9/2^+$	$(56, 4^+_4)$	$1/2$	$N(2220)$	$\Lambda(2350)$	$\Sigma(?)$	$\Xi(?)$	
Decuplet members							
$3/2^+$	$(56, 0^+_0)$	$3/2$	$\Delta(1232)$	$\Sigma(1385)$	$\Xi(1530)$	$\Omega(1672)$	
$3/2^+$	$(56, 0^+_2)$	$3/2$	$\Delta(1600)$	$\Sigma(1690)^{\dagger}$	$\Xi(?)$	$\Omega(?)$	
$1/2^-$	$(70, 1^-_1)$	$1/2$	$\Delta(1620)$	$\Sigma(1750)^{\dagger}$	$\Xi(?)$	$\Omega(?)$	
$3/2^-$	$(70, 1^-_1)$	$1/2$	$\Delta(1700)$	$\Sigma(?)$	$\Xi(?)$	$\Omega(?)$	
$5/2^+$	$(56, 2^+_2)$	$3/2$	$\Delta(1905)$	$\Sigma(?)$	$\Xi(?)$	$\Omega(?)$	
$7/2^+$	$(56, 2^+_2)$	$3/2$	$\Delta(1950)$	$\Sigma(2030)$	$\Xi(?)$	$\Omega(?)$	
$11/2^+$	$(56, 4^+_4)$	$3/2$	$\Delta(2420)$	$\Sigma(?)$	$\Xi(?)$	$\Omega(?)$	

(see below). The mixing is more important for the decay patterns of the states than for their positions. An example are the lowest lying  $(70, 1^-_1)$  states with  $J^P = 1/2^-$  and  $3/2^-$ . The physical states are:

$$|N(1535)1/2^-i = \cos(\Theta_S) |N^2P_M 1/2^-i - \sin(\Theta_S) |N^4P_M 1/2^-i \quad (15.33)$$

$$|N(1520)3/2^-i = \cos(\Theta_D) |N^2P_M 3/2^-i - \sin(\Theta_D) |N^4P_M 3/2^-i \quad (15.34)$$

and the orthogonal combinations for  $N(1650)1/2^-$  and  $N(1700)3/2^-$ . The mixing is large for the  $J^P = 1/2^-$  states ( $\Theta_S \approx -32^\circ$ ), but small for the  $J^P = 3/2^-$  states ( $\Theta_D \approx +6^\circ$ ) [43–45].

All baryons of the ground state multiplets are known. Many of their properties, in particular their masses, are in good agreement even with the most basic versions of the quark model, including harmonic (or linear) confinement and a spin-spin interaction, which is responsible for the octet decuplet mass shifts. A consistent description of the ground-state electroweak properties, however, requires refined relativistic constituent quark models.

**Table 15.8:**  $N$  and  $\Delta$  states in the  $N=0,1,2$  harmonic oscillator bands.  $L^P$  denotes angular momentum and parity,  $S$  the three-quark spin and ‘sym’=A,S,M the symmetry of the spatial wave function. Listed are all possible spin/parity combinations and assignments of experimentally observed states. Only dominant components are indicated. Assignments in the  $N=2$  band are partly tentative.

N	sym	$L^P$	$S$	$N(I = 1/2)$				
2	A	1+	1/2	1/2 <sup>+</sup>	3/2 <sup>+</sup>	-	-	-
2	M	2+	3/2	1/2 <sup>+</sup>	3/2 <sup>+</sup>	5/2 <sup>+</sup>	7/2 <sup>+</sup>	-
2	M	2+	1/2	-	3/2 <sup>+</sup>	5/2 <sup>+</sup>	-	-
2	M	0+	3/2	-	3/2 <sup>+</sup>	-	-	-
2	M	0+	1/2	1/2 <sup>+</sup>	$N(1710)$	-	-	-
2	S	2+	3/2	-	-	-	-	-
2	S	2+	1/2	-	3/2 <sup>+</sup>	$N(1720)$	5/2 <sup>+</sup>	$N(1680)$
2	S	0+	3/2	-	-	-	-	-
2	S	0+	1/2	1/2 <sup>+</sup>	$N(1440)$	-	-	-
1	M	1-	3/2	1/2 <sup>-</sup>	$N(1650)$	3/2 <sup>-</sup>	$N(1700)$	5/2 <sup>-</sup>
1	M	1-	1/2	1/2 <sup>-</sup>	$N(1535)$	3/2 <sup>-</sup>	$N(1520)$	-
0	S	0+	3/2	-	-	-	-	-
0	S	0+	1/2	1/2 <sup>+</sup>	$N(938)$	-	-	-
N	sym	$L^P$	$S$	$\Delta(I = 3/2)$				
2	A	1+	1/2	-	-	-	-	-



## ENTRI

2	M	2+	3/2	-	-	-	-
2	M	2+	1/2	-	3/2 <sup>+</sup>	5/2 <sup>+</sup>	-
2	M	0+	3/2	-	-	-	-
2	M	0+	1/2	1/2 <sup>+</sup> $\Delta(1750)$	-	-	-
2	S	2+	3/2	1/2 <sup>+</sup> $\Delta(1910)$	3/2 <sup>+</sup> $\Delta(1920)$	5/2 <sup>+</sup> $\Delta(1905)$	7/2 <sup>+</sup> $\Delta(1950)$
2	S	2+	1/2	-	-	-	-
2	S	0+	3/2	-	3/2 <sup>+</sup> $\Delta(1600)$	-	-
2	S	0+	1/2	-	-	-	-
1	M	1-	3/2	-	-	-	-
1	M	1-	1/2	1/2 <sup>-</sup> $\Delta(1620)$	3/2 <sup>-</sup> $\Delta(1700)$	-	-
0	S	0+	3/2	-	3/2 <sup>+</sup> $\Delta(1232)$	-	-
0	S	0+	1/2	-	-	-	-

The situation for the excited states is much less clear. The assignment of some experimentally observed states with strange quarks to model configurations is only tentative and in many cases candidates are completely missing. Melde, Plessas and Sengl [42] have calculated baryon properties in relativistic constituent quark models, using one-gluon exchange and Goldstone-boson exchange for the modeling of the hyperfine interactions (see Sec. 15.5 on Dynamics). Both types of models give qualitatively comparable results, and underestimate in general experimentally observed decay widths. Nevertheless, in particular on the basis of the observed decay patterns, the authors have assigned some additional states with strangeness to the SU(3) multiplets and suggest re-assignments for a few others. Among the new assignments are states with weak experimental evidence (two or three star ratings) and partly without firm spin/parity assignments, so that further experimental efforts are necessary before final conclusions can be drawn. We have added their suggestions in Table 15.7.




Figure 15.6: Excitation spectrum of the nucleon. Compared are the positions of the excited states identified in experiment, to those predicted by a relativized quark model calculation. Left hand side: isospin  $I = 1/2$   $N$ -states, right hand side: isospin  $I = 3/2$   $\Delta$ -states. Experimental: (columns labeled 'exp'), three- and four-star states are indicated by full lines (two-star dashed lines, one-star dotted lines). At the very left and right of the figure, the spectroscopic notation of these states is given. Quark model [46,47]: (columns labeled 'QM'), all states for the  $N=1,2$  bands, low-lying states for the  $N=3,4,5$  bands. Full lines: at least tentative assignment to observed states, dashed lines: so far no observed counterparts. Many of the assignments between predicted and observed states are highly tentative.

In the non-strange sector there are two main problems which are illustrated in Fig. 15.6, where the experimentally observed excitation spectrum of the nucleon ( $N$  and  $\Delta$  resonances) is compared to the results of a typical quark model calculation [46]. The lowest states from the  $N=2$  band, the  $N(1440)1/2^+$ , and the  $\Delta(1600)3/2^+$ , appear lower than the negative parity states from the  $N=1$  band (see Table 15.8) and much lower than predicted by most models. Also negative parity  $\Delta$  states from the  $N=3$  band ( $\Delta(1900)1/2^-$ ,  $\Delta(1940)3/2^-$ , and  $\Delta(1930)5/2^-$ ) are too low in energy. Part of the problem could be experimental. Among the negative parity  $\Delta$  states, only the  $\Delta(1930)5/2^-$  has three stars and the uncertainty in the position of the  $\Delta(1600)3/2^+$  is large (1550 - 1700 MeV).

Furthermore, many more states are predicted than observed. This has been known for a long time as the 'missing resonance' problem [43]. Up to an excitation energy of 2.4 GeV, about 45  $N$  states are predicted, but only 20 are established (four- or three-star; see Note on  $N$  and  $\Delta$  Resonances for the rating of the status of resonances) and 5 are tentative (two- or one-star). Even for the  $N=2$  band, up to now only half of the predicted states have been observed. However, there is some recent progress. The total

number of states has not much changed but the number of states with four- or three-star rating has increased from 14 to 20 compared to the 2018 PDG particle listings. Most of this progress is due to the programs concentrating on the study of meson photoproduction reactions, while the most recent partial wave analysis of elastic pion scattering and charge exchange data by Arndt and collaborators [48] found no evidence for almost half of the states listed in this review (and included in Fig. 15.6). Such analyses are of course biased against resonances which couple only weakly to the  $N\pi$  channel. Quark model predictions for the couplings to other hadronic channels and to photons are given in Ref. [46]. The large experimental effort ongoing at several electron accelerators to study the baryon resonance spectrum with real and virtual photon-induced meson production reactions includes the search for as-yet-unobserved states, as well as detailed studies of the properties of the low lying states (decay patterns, electromagnetic couplings, magnetic moments, *etc.*) (see Ref. [49] for reviews). There are two major new aspects of this program. The investigation of single and double polarization observables allows via the study of interference terms access to small partial waves that do not leave a footprint in unpolarized cross sections. An example for the impact of such data is given by a comparison of results from different multipole analyses of pion photoproduction [50]. It shows clearly that with the inclusion of polarization observables the reaction model results start to converge. This will in the near future much improve the data basis for excited baryons in the light quark sector.

The other aspect is the study of final states with meson pairs, in particular  $\pi\pi$  and  $\pi\eta$  pairs, which made large progress during the last few years. This is important for higher lying states, which in the quark model may have both possible oscillations excited. Such states can be expected to decay in sequential processes de-exciting the two oscillations step-by-step so that they couple strongly to multiple-meson final states but not to single-meson production. Detailed analyses of such data are for example given in [51, 52] and had already significant impact on partial wave analyses.

In quark models, the number of excited states is determined by the effective degrees of freedom, while their ordering and decay properties are related to the residual quark - quark interaction. An overview of quark models for baryons is given in [53], recent discussions of baryon spectroscopy are given in [34,41]. The effective degrees of freedom in the standard nonrelativistic quark model are three equivalent valence quarks with one-gluon exchange-motivated, flavor-independent colormagnetic interactions. The QCD aspect of gluon-gluon interactions is emphasized by the hypercentral quark model [54,55], which includes in a natural way three-body forces between the quarks. A different class of models uses interactions which give rise to a quark - diquark clustering of the baryons: for a review see [56]. If there is a tightly bound diquark, only two degrees of freedom are available at low energies, and thus *fewer* states are predicted. Furthermore, selection rules in the decay pattern may arise from the quantum numbers of the diquark. *More* states are predicted by collective models of the baryon like the algebraic approach in [57]. In this approach, the quantum numbers of the valence quarks are distributed over a Y-shaped string-like configuration, and additional states arise *e.g.*, from vibrations of the strings. *More* states are also predicted in the framework of flux-tube models, see [58], which are motivated by lattice QCD. In addition to the quark degrees of freedom, flux-tubes responsible for the confinement of the quarks are considered as degrees of freedom. These models include hybrid baryons containing explicit excitations of the gluon fields. However, since all half integral  $J^P$  quantum numbers are possible for ordinary baryons, such 'exotics' will be very hard to identify, and probably always mix with ordinary states. So far, the experimentally observed number of states is still far lower even than predicted by the quark-diquark models.

The influence of chiral symmetry on the excitation spectrum of the nucleon has been debated from a somewhat different perspective. Chiral symmetry, the fundamental symmetry of QCD, is strongly broken for the low lying states, resulting in large mass differences of parity partners like the  $J^P = 1/2^+ N(938) 1/2^+$

ground state and the  $J^P = 1/2^- N(1535) 1/2^-$  excitation. However, at higher excitation energies there is some evidence for parity doublets and even some very tentative suggestions for full chiral multiplets of  $N^7$  and  $\Delta$  resonances. An effective restoration of chiral symmetry at high excitation energies due to a decoupling from the quark condensate of the vacuum has been discussed (see Ref. [59] for recent reviews) as a possible cause. In this case, the mass generating mechanisms for low and high lying states would be essentially different. As a further consequence, the parity doublets would decouple from pions, so that experimental bias would be worse. However, parity doublets might also arise from the spin-orbital dynamics of the 3-quark system. Presently, the status of data does not allow final conclusions.

The most recent developments on the theory side are the first unquenched lattice calculations for the excitation spectrum discussed in Sec 15.6. The results are basically consistent with the level counting of  $SU(6) \otimes O(3)$  in the standard non-relativistic quark model and show no indication for quark-diquark structures or parity doubling. Consequently, there is as yet no indication from lattice that the mis-match between the excitation spectrum predicted by the standard quark model and experimental observations is due to inappropriate degrees of freedom in the quark model.

### 15.5 Dynamics

Quantum chromodynamics (QCD) is well-established as the theory for the strong interactions. As such, one of the goals of QCD is to predict the spectrum of strongly-interacting particles. To date, the only first-principles calculations of spectroscopy from QCD use lattice methods. These are the subject of Sec. 15.6. These calculations are difficult and unwieldy, and many interesting questions do not have a good lattice-based method of solution. Therefore, it is natural to build models, whose ingredients are abstracted from QCD, or from the low-energy limit of QCD (such as chiral Lagrangians) or from the data itself. The words “quark model” are a shorthand for such phenomenological models. Many specific quark models exist, but most contain a similar basic set of dynamical ingredients. These include:

1. A confining interaction, which is generally spin-independent (*e.g.*, harmonic oscillator or linear confinement);
2. Different types of spin-dependent interactions:
  - a) commonly used is a color-magnetic flavor-independent interaction modeled after the effects of gluon exchange in QCD (see *e.g.*, Ref. [60]). For example, in the *S*-wave states, there is a spin-spin hyperfine interaction of the form

$$H_{HF} = -\alpha_s M \sum_{i>j} (\vec{\sigma}_i \cdot \vec{\lambda}_a)_i (\vec{\sigma}_j \cdot \vec{\lambda}_a)_j, \quad (15.35)$$

where  $M$  is a constant with units of energy,  $\lambda_a$  ( $a = 1, \dots, 8$ ) is the set of  $SU(3)$  unitary spin matrices, defined in the review “ $SU(3)$  Isoscalar Factors and Representation Matrices,” and the sum runs over constituent quarks or antiquarks. Spin-orbit interactions, although allowed, seem to be small in general, but a tensor term is responsible for the mixing of states with the same  $J^P$  but different  $L, S$  combinations.

- b) other approaches include flavor-dependent short-range quark forces from instanton effects (see *e.g.*, [61,62]). This interaction acts only on scalar, isoscalar pairs of quarks in a relative *S*-wave state:

$$h q^2; S, L, T / W / q^2; S, L, T i = -4g \delta_{S,0} \delta_{L,0} \delta_{T,0} W \quad (15.36)$$

where  $W$  is the radial matrix element of the contact interaction.

c) a rather different and controversially discussed approach is based on flavor-dependent spin-spin forces arising from one-boson exchange. The interaction term is of the form:

$$H_{HF} \propto \sum_{i < j} V(\sim r_{ij}) \lambda_i^F \cdot \lambda_j^F \sim \sigma_i \cdot \sigma_j \quad (15.37)$$

where the  $\lambda_i^F$  are in flavor space (see *e.g.*, Ref. [63]).

3. A strange quark mass somewhat larger than the up and down quark masses, in order to split the SU(3) multiplets;
4. In the case of spin-spin interactions (iia,c), a flavor-symmetric interaction for mixing  $qq^-$  configurations of different flavors (*e.g.*,  $uu^- \leftrightarrow dd^- \leftrightarrow ss^-$ ), in isoscalar channels, so as to reproduce *e.g.*, the  $\eta - \eta^0$  and  $\omega - \varphi$  mesons.

These ingredients provide the basic mechanisms that determine the hadron spectrum in the standard quark model.

## 15.6 Lattice Calculations of Hadronic Spectroscopy

Lattice calculations are a major source of information about QCD masses and matrix elements. The necessary theoretical background is given in Sec. 17 of this *Review*. Here we confine ourselves to some general comments and illustrations of lattice calculations for spectroscopy.

In general, the cleanest lattice results come from computations of processes in which there is only one particle in the simulation volume. These quantities include masses of hadrons, simple decay constants, like pseudoscalar meson decay constants, and semileptonic form factors (such as the ones appropriate to  $B \rightarrow D\ell\nu$ ,  $K\ell\nu$ ,  $\pi\ell\nu$ ). The cleanest predictions for masses are for states which have narrow decay widths and are far below any thresholds to open channels, since the effects of final state interactions are not yet under complete control on the lattice. As a simple corollary, the lightest state in a channel is easier to study than the heavier ones. “Difficult” states for the quark model (such as exotics) are also difficult for the lattice because of the lack of simple operators which couple well to them.

Good-quality modern lattice calculations will present multi-part error budgets with their predictions. A small part of the uncertainty is statistical, from sample size. Typically, the quoted statistical uncertainty includes uncertainty from a fit: it is rare that a simulation computes one global quantity which is the desired observable. Simulations which include virtual quark-antiquark pairs (also known as “dynamical quarks” or “sea quarks”) are often done at up and down quark mass values heavier than the experimental ones, and it is then necessary to extrapolate in these quark masses. Simulations can work at the physical values of the heavier quarks’ masses. They are always done at nonzero lattice spacing, and so it is necessary to extrapolate to zero lattice spacing. Some theoretical input is needed to do this. Much of the uncertainty in these extrapolations is systematic, from the choice of fitting function. Other systematics include the effect of finite simulation volume, the number of flavors of dynamical quarks actually simulated, and technical issues with how these dynamical quarks are included. The particular choice of a fiducial mass (to normalize other predictions) is not standardized; there are many possible choices, each with its own set of strengths and weaknesses, and determining it usually requires a second lattice simulation from that used to calculate the quantity under consideration.

A systematic error of major historical interest is the “quenched approximation,” in which dynamical quarks are simply left out of the simulation. This was done because the addition of these virtual pairs presented an expensive computational problem. No generally-accepted methodology has ever allowed

one to correct for quenching effects, short of redoing all calculations with dynamical quarks. Recent advances in algorithms and computer hardware have rendered it obsolete.

With these brief remarks, we turn to examples. The field of lattice QCD simulations is vast, and so it is not possible to give a comprehensive review of them in a small space. The history of lattice QCD simulations is a story of thirty years of incremental improvements in physical understanding, algorithm development, and ever faster computers, which have combined to bring the field to a present state where it is possible to carry out very high quality calculations. We present a few representative illustrations, to show the current state of the art.

By far, the major part of all lattice spectroscopy is concerned with that of the light hadrons, and so we illustrate results in Fig. 15.7, a comprehensive summary provided by A. Kronfeld (private communication; see also [64]).




Figure 15.7: Hadron spectrum from lattice QCD. Comprehensive results for mesons and baryons are from MILC [65,66], PACS-CS [67], BMW [68], QCDSF [69], and ETM [70]. Results for  $\eta$  and  $\eta'$  are from RBC & UKQCD [10], Hadron Spectrum [71] (also the only  $\omega$  mass), UKQCD [72], and Michael, Ottnad, and Urbach [73]. Results for heavy-light hadrons from Fermilab-MILC [74], HPQCD [75,76], and Mohler and Woloshyn [77]. Circles, squares, diamonds, and triangles stand for staggered, Wilson, twisted-mass Wilson, and chiral sea quarks, respectively. Asterisks represent anisotropic lattices. Open symbols denote the masses used to fix parameters. Filled symbols (and asterisks) denote results. Red, orange, yellow, green, and blue stand for increasing numbers of ensembles (i.e., lattice spacing and sea quark mass) Black symbols stand for results with 2+1+1 flavors of sea quarks. Horizontal bars (gray boxes) denote experimentally measured masses (widths).  $b$ -flavored meson masses are offset by  $-4000$  MeV.

Flavor singlet mesons are at the frontier of lattice QCD calculations, because one must include the effects of “annihilation graphs,” for the valence  $q$  and  $q^-$ . Recently, several groups, Refs. [10,72, 78], have reported calculations of the  $\eta$  and  $\eta'$  mesons. The numbers of [10] are typical, finding masses of 573(6) and 947(142) MeV for the  $\eta$  and  $\eta'$ . The singlet-octet mixing angle (in the conventions of Table 15.2) is  $\vartheta_{lin} = -14.1(2.8)^\circ$ .

The spectroscopy of mesons containing heavy quarks has become a truly high-precision endeavor. These simulations use Non-Relativistic QCD (NRQCD) or Heavy Quark Effective Theory (HQET), systematic expansions of the QCD Lagrangian in powers of the heavy quark velocity, or the heavy quark mass. Terms in the Lagrangian have obvious quark model analogs, but are derived directly from QCD. For example, the heavy quark potential is a derived quantity, extracted from simulations. Fig. 15.8 shows the mass spectrum for mesons containing at least one heavy ( $b$  or  $c$ ) quark from Ref. [76]. It also contains results from Ref. [79,80]. The calculations uses a discretization of nonrelativistic QCD for bottom quarks with charm and lighter quarks being handled with an improved relativistic action. Four flavors ( $u, d, s, c$ ) of dynamical quarks are included.




Figure 15.8: Spectroscopy for mesonic systems containing one or more heavy quarks (adapted from Ref. [76]. Particles whose masses are used to fix lattice parameters are shown with crosses; the authors distinguish between “predictions” and “postdictions” of their calculation. Lines represent experiment.

Fig. 15.9 shows a compilation of recent lattice results for doubly and triply charmed baryons, provided by S. Meinel [81]. The state recently announced by LHCb [36] is also shown. Note that the lattice calculations for the mass of this state were predictions, not postdictions.

Recall  
lattice

that




Figure 15.9: Comparison of lattice QCD results for the doubly and triply charmed baryon masses. Labels are Liu, *et al.*, [82]; Briceño, *et al.*, [83]; Namekawa, *et al.*, [84]; Padmanath, *et al.*, [85]; Alexandrou, *et al.*, [70]; Brown, *et al.*, [86]; Perez-Rubio *et al.*, [87]; Alexandrou and Kallidonis 2017, [88]. Only calculations with dynamical light quarks are included; for the doubly charmed baryons, only calculations were performed at or extrapolated to the physical pion mass are shown. Results without estimates of systematic uncertainties are labeled “stat. only”. The lattice spacing values used in the calculations are also given;  $a = 0$  indicates that the results have been extrapolated to the continuum limit. In the plot of the doubly charmed baryons, the recently announced experimental result for the  $\Xi_{cc}^+$  mass from LHCb [36] is shown with a horizontal line.

bers appropriate to the desired states, compute correlation functions of these operators, and fit the correlation functions to functional forms parametrized by a set of masses and matrix elements. As we move away from hadrons which can be created by the simplest quark model operators (appropriate to the lightest meson and baryon multiplets) we encounter a host of new problems: either no good interpolating fields, or too many possible interpolating fields, and many states with the same quantum numbers. Techniques for dealing with these interrelated problems vary from collaboration to collaboration, but all share common features: typically, correlation functions from many different interpolating fields are used, and the signal is extracted in what amounts to a variational calculation using the chosen operator basis. In addition to mass spectra, wave function information can be garnered from

Most hadrons are resonances, and lattice calculations will have to deal with this fact as the




Figure 15.11: Spin-identified spectrum of nucleons and deltas, from lattices where  $m_\pi = 396$  MeV, in units of the calculated  $\Omega$  mass, from Ref. [90]. The colors just correspond to the different  $J$  assignments: grey for  $J = 1/2$ , red for  $J = 3/2$ , green for  $5/2$ , blue for  $7/2$ .

quark masses are taken ever smaller. The actual calculation is of the combined mass of two (or more) hadrons in a box of finite size. The combined mass is shifted from being the sum of the individual masses because the finite box forces the hadrons to interact with each other. The volume-dependent mass shift yields the phase shift for the continuum scattering amplitude, which in turn can be used to extract the resonance mass and width, with some degree of modeling. So far only two-body resonances, the rho meson and a few others, have been well studied. This is an active research area. A recent review, [91], summarizes the situation, and example of a calculation of the rho meson decay width is [92]. The mass and decay width of the  $f_0(500)$  have recently been computed in [93]. Ref. [94] studies the decay width of the  $\Delta(1238)$ . Lattice calculations relevant to the extra states observed in the charmonium spectrum (Sec. 15.3) are difficult, because the states sit high in the spectrum of most channels and due to the number of nearby multiparticle states.

In Fig. 15.4 we showed a figure from [18] presenting a lattice prediction for the glueball mass spectrum in quenched approximation. A true QCD prediction of the glueball spectrum requires dynamical light quarks and (because glueball operators are intrinsically noisy) high statistics. Only recently have the first useful such calculations appeared, in [95, 96]. Fig. 15.12 shows results from [95], done with dynamical  $u$ ,  $d$  and  $s$  quarks at two lattice spacings, 0.123 and 0.092 fm, along with comparisons to the quenched lattice calculation of [17] and to experimental isosinglet mesons. The dynamical simulation is, of course, not the last word on this subject, but it shows that the effects of quenching seem to be small.




Figure 15.12: Lattice QCD predictions for glueball masses. The open and closed circles are the larger and smaller lattice spacing data of the full QCD calculation of glueball masses of Ref. [95]. Squares are the quenched data for glueball masses of Ref. [17]. The bursts labeled by particle names are experimental states with the appropriate quantum numbers.

As a final part of spectroscopy we mention electromagnetic mass splittings (such as the neutron - proton mass difference). They are interesting but difficult. These calculations are important for determining the values of the quark masses (for a discussion see the review in the PDG). Knowing that the neutron is heavier than the proton tells us that these splittings have a complicated origin. One part of the shift is because the up and down quarks have slightly different masses. The second is that the quarks have (different) charges. Phenomenologists (compare Ref. [97]) combine Coulomb forces and spin-dependent electromagnetic hyperfine interactions to model their charge effects. In order to compute hadronic mass differences on the lattice, electromagnetic interactions must be included in the simulations. This creates a host of technical issues. An important one is that electromagnetic interactions are long range, but lattice simulations are done in finite volumes. The theoretical situation is summarized in the recent review [98]. A recent calculation, Ref. [99], has presented the first results for electromagnetic mass splittings in the baryon octet, with good agreement with observation. Ref. [100] has calculations for meson splittings.