


# Legendre Polynomials and Functions

## Outline

<b>Background and Definitions</b> .....	2
<b>Definitions</b> .....	3
<b>Theory</b> .....	4
<b>Legendre's Equation, Functions and Polynomials</b> .....	4
<b>Legendre's Associated Equation and Functions</b> .....	13
<b>Assigned Problems</b> .....	16



## Background and Definitions

The ordinary differential equation referred to as *Legendre's differential equation* is frequently encountered in physics and engineering. In particular, it occurs when solving Laplace's equation in spherical coordinates.

Adrien-Marie Legendre (September 18, 1752 - January 10, 1833) began using, what are now referred to as Legendre polynomials in 1784 while studying the attraction of spheroids and ellipsoids. His work was important for geodesy.

### 1. Legendre's Equation and Legendre Functions

The second order differential equation given as

$$(1 - x^2) \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + n(n + 1)y = 0 \quad n > 0, \quad |x| < 1$$

is known as Legendre's equation. The general solution to this equation is given as a function of two Legendre functions as follows

$$y = AP_n(x) + BQ_n(x) \quad |x| < 1$$

where

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x - 1)^n \quad \text{Legendre function of the first kind}$$

$$Q_n(x) = -\frac{1}{2} P_n(x) \ln \frac{1+x}{1-x} \quad \text{Legendre function of the second kind}$$

### 2. Legendre's Associated Differential Equation

Legendre's associated differential equation is given as

$$(1 - x^2) \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + \left[ n(n + 1) - \frac{m^2}{1 - x^2} \right] y = 0$$

If we set  $m = 0$  in this equation the differential equation reduces to Legendre's equation.

The general solution to Legendre's associated equation is given as

$$y = A P_n^m(x) + B Q_n^m(x)$$

where  $P_n^m(x)$  and  $Q_n^m(x)$  are called the associated Legendre functions of the first and second kind given as

$$P_n^m(x) = (1-x^2)^{m/2} \frac{d^m P_n(x)}{dx^m}$$

$$Q_n^m(x) = (1-x^2)^{m/2} \frac{d^m Q_n(x)}{dx^m}$$



## Legendre's Equation and Its Solutions

Legendre's differential equations is

$$(1-x^2) \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + n(n+1)y = 0 \quad n > 0, |x| < 1$$

or equivalently

$$\frac{d}{dx} \left[ (1-x^2) \frac{dy}{dx} \right] + n(n+1)y = 0 \quad n > 0, |x| < 1$$

Solutions of this equation are called Legendre functions of order  $n$ . The general solution can be expressed as

$$y = AP_n(x) + BQ_n(x) \quad |x| < 1$$

where  $P_n(x)$  and  $Q_n(x)$  are *Legendre Functions* of the first and second kind of order  $n$ .

If  $n = 0, 1, 2, 3, \dots$  the  $P_n(x)$  functions are called *Legendre Polynomials* of order  $n$  and are given by Rodrigue's formula.

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$$

Legendre functions of the first kind ( $P_n(x)$ ) and second kind ( $Q_n(x)$ ) of order  $n = 0, 1, 2, 3$  are shown in the following two plots

The first several Legendre polynomials are listed below

$$P_0(x) = 1$$

$$P_3(x) = \frac{1}{8}(5x^3 - 3x)$$

$$P_1(x) = x$$

$$P_3(x) = \frac{1}{8}(35x^3 - 30x^2 + 3x)$$

$$P_2(x) = \frac{1}{2}(3x^2 - 1)$$

$$P_3(x) = \frac{1}{8}(63x^3 - 70x^2 + 15x)$$

The recurrence formula is

$$P_{n+1}(x) = \frac{2n+1}{n+1}xP_n(x) - \frac{n}{n+1}P_{n-1}(x)$$

$$P_{n+1}(x) - P_{n-1}(x) = (2n+2)P_n(x)$$

can be used to obtain higher order polynomials. In all cases  $P_n(1) = 1$  and  $P_n(-1) = (-1)^n$

## Orthogonality of Legendre Polynomials

The Legendre polynomials  $P_m(x)$  and  $P_n(x)$  are said to be orthogonal in the interval  $-1 \leq x \leq 1$  provided

$$\int_{-1}^1 P_m(x) P_n(x) dx = 0 \quad m \neq n$$

and as a result we have

$$\int_{-1}^1 [P_n(x)]^2 dx = \frac{2}{2n+1} \quad m = n$$




Figure 5.1: Legendre function of the first kind,  $P_n(x)$




Figure 5.2: Legendre function of the second kind,  $Q_n(x)$

### Orthogonal Series of Legendre Polynomials

Any function  $f(x)$  which is finite and single-valued in the interval  $-1 \leq x \leq 1$ , and which has a finite number or discontinuities within this interval can be expressed as a series of Legendre polynomials.

We let

$$f(x) = A_0P_0(x) + A_1P_1(x) + A_2P_2(x) + \dots \quad -1 \leq x \leq 1$$

$$= \sum_{n=0}^{\infty} A_n P_n(x)$$

Multiplying both sides by  $P_m(x) dx$  and integrating with respect to  $x$  from  $x = -1$  to  $x = 1$  gives

$$\int_{-1}^1 f(x)P_m(x) dx = \sum_{n=0}^{\infty} A_n \int_{-1}^1 P_m(x)P_n(x) dx$$

By means of the orthogonality property of the Legendre polynomials we can write

$$A_n = \frac{n+1}{2} \int_{-1}^1 f(x)P_n(x) dx \quad n = 0, 1, 2, 3, \dots$$

Since  $P_n(x)$  is an even function of  $x$  when  $n$  is even, and an odd function when  $n$  is odd, it follows that if  $f(x)$  is an even function of  $x$  the coefficients  $A_n$  will vanish when  $n$  is odd; whereas if  $f(x)$  is an odd function of  $x$ , the coefficients  $A_n$  will vanish when  $n$  is even. Thus for an even function  $f(x)$  we have

$$A_n = \begin{cases} 0 & n \text{ is odd} \\ (2n+1) \int_0^1 f(x)P_n(x) dx & n \text{ is even} \end{cases}$$

whereas for an odd function  $f(x)$  we have

$$= \begin{cases} (2n+1) \int_0^1 A_n f(x)P_n(x) dx & n \text{ is odd} \\ 0 & n \text{ is even} \end{cases}$$

When  $x = \cos\theta$  the function  $f(\theta)$  can be written

$$f(\theta) = \sum_{n=0}^{\infty} A_n P_n(\cos\theta) \quad 0 \leq \theta \leq \pi$$

where

$$= \frac{n+1}{2} \int_0^\pi f(\theta) P_n(\cos\theta) \sin\theta \, d\theta \quad n = 0, 1, 2, 3, \dots$$

### Some Special Results Legendre Polynomials

Integral form

$$P_n(x) = \frac{1}{\pi} \int_0^\pi \frac{x + \sqrt{x^2 - 1} \cos t}{2} P_n(\cos t) \, dt$$

Values of  $P_n(x)$  at  $x = 0$  and  $x = \pm 1$

$$P_{2n}(0) = \frac{(-1)^n \Gamma(n + 1/2)}{\sqrt{\pi} \Gamma(n + 1)} \quad P_{2n+1}(0) = 0$$

$$P_{2n}(0) = 0 \quad P_{2n+1}(0) = \frac{(-1)^n 2\Gamma(n + 3/2)}{\sqrt{\pi} \Gamma(n + 1)}$$

$$P_n(1) = 1 \quad P_n(-1) = (-1)^n$$

$$P_n(1) = \frac{n(n+1)}{2} \quad P_n(-1) = (-1)^{n-1} \frac{n(n+1)}{2}$$

$$|P_n(x)| \leq 1$$

The primes denote differentiation with respect to  $x$  therefore

$$P_n'(1) = \frac{dP_n(x)}{dx} \text{ at } x = 1$$

### Generating Function for Legendre Polynomials

If  $A$  is a fixed point with coordinates  $(x_1, y_1, z_1)$  and  $P$  is the variable point  $(x, y, z)$  and the distance  $AP$  is denoted by  $R$ , we have



$$R^2 = (x - x_1)^2 + (y - y_1)^2 + (z - z_1)^2$$

From the theory of Newtonian potential we know that the potential at the point P due to a unit mass situated at the point A is given by

$$\phi = \frac{C}{R}$$

where C is some constant. It can be shown that this function is a solution of Laplace's equation.

In some circumstances, it is desirable to expand  $\phi$  in powers of  $r$  or  $r^{-1}$  where  $r =$

$\sqrt{x^2 + y^2 + z^2}$  is the distance from the origin O to the point P.

z




Figure 5.3: Generating Function for Legendre Polynomials

$$a = |\vec{OA}|$$

$$r = |\vec{OB}|$$

$$\phi = \frac{C}{R} = \frac{C}{\sqrt{r^2 + a^2 - 2ra \cos \theta}} = \frac{C}{r} \frac{1}{\sqrt{1 + \frac{a^2}{r^2} - 2\frac{a}{r} \cos \theta}}$$

Through substitution we can write

$$C \phi = \frac{[1 - 2xt + t^2]^{-1/2}}{r}$$

where

$$t = \frac{a}{r}, \quad x = \cos \theta$$

Therefore

$$C \phi \equiv \frac{g(x,t)}{r}$$

We introduce the angle  $\theta$  between the vectors  $OA$  and  $OP$  and write

$$R^2 = r^2 + a^2 - 2ra \cos \theta$$

where  $a = |OA|$ . If we let  $r/R = t$  and  $x = \cos \theta$ , then

$$g(x,t) = (1 - 2xt + t^2)^{-1/2}$$

is defined as the generating function for  $P_n(x)$ . Expanding by the binomial expansion we have

$$g(x,t) = \sum_{n=0}^{\infty} \frac{1}{2^n} \binom{-1/2}{n} (2xt - t^2)^n$$

where the symbol  $(\alpha)_n$  is defined by

$$(\alpha)_n = \alpha(\alpha + 1)(\alpha + 2) \dots (\alpha + n - 1) = \prod_{k=0}^{n-1} (\alpha + k)$$

$$(\alpha)_0 = 1$$

$(\alpha)_n$  is referred to as the Pochhammer symbol and  $(\alpha, n)$  is the Appel's symbol.

Thus we have

$$g(x, t) = \sum_{n=0}^{\infty} \frac{(1/2)_n}{n!} \sum_{k=0}^n \frac{(2x)^{n-k} t^{n-k} (-t^2)^k}{k!(n-k)!}$$

which can be written as

$$g(x, t) = (1 - 2xt + t^2)^{-1/2} = \sum_{n=0}^{\infty} \left[ \sum_{k=0}^{n/2} \frac{(-1)^k (2n - 2k)! x^{n-2k}}{2^k k!(n-2k)!(n-k)!} \right] t^n$$

The coefficient of  $t^n$  is the Legendre polynomial  $P_n(x)$ , therefore

$$g(x, t) = (1 - 2xt + t^2)^{-1/2} = \sum_{n=0}^{\infty} P_n(x) t^n \quad |x| \leq 1, |t| < 1$$

### Legendre Functions of the Second Kind

A second and linearly independent solution of Legendre's equation for  $n$ =positive integers are called Legendre functions of the second kind and are defined by

$$Q_n(x) = \frac{1}{2} P_n(x) \ln \frac{1+x}{1-x} + W_{n-1}(x)$$

where

$n$

**E ▶ ENTRI**

$$Q_{n-1}(x) = \sum_{m=1}^n \frac{1}{m} P_{m-1}(x) P_{n-m}(x) W_{nm}(x)$$

is a polynomial of the  $(n-1)$  degree. The first term of  $Q_n(x)$  has logarithmic singularities at  $x = \pm 1$  or  $\theta = 0$  and  $\pi$ .

The first few polynomials are listed below

$$Q_0(x) = \frac{1}{2} \ln \frac{1+x}{1-x}$$

$$Q_1(x) = P_1(x)Q_0(x) - 1$$

$$Q_2(x) = P_2(x)Q_0(x) - \frac{3}{2}x$$

$$Q_3(x) = P_3(x)Q_0(x) - \frac{5}{2}x^2 + \frac{2}{3}$$

showing the even order functions to be odd in  $x$  and conversely.

The higher order polynomials  $Q_n(x)$  can be obtained by means of recurrence formulas exactly analogous to those for  $P_n(x)$ .

Numerous relations involving the Legendre functions can be derived by means of complex variable theory. One such relation is an integral relation of  $Q_n(x)$

$$Q_n(x) = \int_0^\infty \left[ x + \sqrt{x^2 - 1} \cosh \theta \right]^{-n-1} d\theta \quad |x| > 1$$

and its generating function

$$(1 - 2xt + t^2)^{-1/2} \cosh^{-1} \frac{t-x}{\sqrt{x^2-1}} = \sum_{n=0}^{\infty} Q_n(x)t^n$$

## Some Special Values of $Q_n(x)$

$$Q_{2n+1}(0) = 0 \qquad Q_{2n}(0) = (-1)^{n+1} \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdots (2n-1)}$$

$$Q_n(1) = \infty \qquad Q_n(-x) = (-1)^{n+1} Q_n(x)$$

## Legendre's Associated Differential Equation

The differential equation

$$(1-x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + \left[ n(n+1) - \frac{m^2}{1-x^2} \right] y = 0$$

is called Legendre's associated differential equation. If  $m = 0$ , it reduces to Legendre's equation. Solutions of the above equation are called associated Legendre functions. We will restrict our discussion to the important case where  $m$  and  $n$  are non-negative integers. In this case the general solution can be written

$$y = A P_n^m(x) + B Q_n^m(x)$$

where  $P_n^m(x)$  and  $Q_n^m(x)$  are called the associated Legendre functions of the first and second kind respectively. They are given in terms of ordinary Legendre functions.

$$P_n^m(x) = (1-x^2)^{m/2} \frac{d^m P_n(x)}{dx^m}$$

$$Q_n^m(x) = (1-x^2)^{m/2} \frac{d^m Q_n(x)}{dx^m}$$

The  $P_n^m(x)$  functions are bounded within the interval  $-1 \leq x \leq 1$  whereas  $Q_n^m(x)$  functions are unbounded at  $x = \pm 1$ .

## Special Associated Legendre Functions of the First Kind

$$P_n^0(x) = P_n(x)$$

$$P_n(x) = \frac{1}{n!} \frac{d^n}{dx^n} (x^2 - 1)^n = 0 \quad m > n$$

$$P_n(x) = \frac{1}{n!} \frac{d^n}{dx^n} (x^2 - 1)^n$$

$$P_1(x) = (1 - x^2)^{1/2}$$

$$P_2(x) = 3x(1 - x^2)^{1/2}$$

$$P_2^2(x) = 3(1 - x^2)$$

$$P_3(x) = \frac{3}{2}(5x^2 - 1)(1 - x^2)^{1/2}$$

$$P_3^2(x) = 15x(1 - x^2)$$

$$P_3^3(x) = 15(1 - x^2)^{3/2}$$

Other associated Legendre functions can be obtained by the recurrence formulas.

### Recurrence Formulas for $P_n^m(x)$

$$(n + 1 - m)P_{n+1}^m(x) = (2n + 1)xP_n^m(x) - (n + m)P_{n-1}^m(x)$$

$$P_{n+2}^m(x) = \frac{2(m + 1)}{(1 - x^2)^{1/2}} x P_{n+1}^m - (n - m)(n + m + 1)P_n^m(x)$$

### Orthogonality of $P_n^m(x)$

As in the case of Legendre polynomials, the Legendre functions  $P_n^m(x)$  are orthogonal in the interval  $-1 \leq x \leq 1$

$$\int_{-1}^1 P_n^m(x) P_k^m(x) dx = 0 \quad n \neq k$$

and also

$$\int_{-1}^1 [P_n^m(x)]^2 dx = \frac{2}{2n+1} \frac{(n+m)!}{(n-m)!}$$

## Orthogonality Series of Associated Legendre Functions

Any function  $f(x)$  which is finite and single-valued in the interval  $-1 \leq x \leq 1$  can be expressed as a series of associated Legendre functions

$f(x) = A_m P_{m1}(x) + A_{m+1} P_{m+1}(x) + A_{m+2} P_{m+2}(x) + \dots$   
 where the coefficients are determined by means of

$$A_k = \frac{k+1}{2} \frac{(k-m)!}{(k+m)!} \int_{-1}^1 f(x) P_k^m(x) dx$$

## Assigned Problems

### Problem Set for Legendre Functions and Polynomials

1. Obtain the Legendre polynomial  $P_4(x)$  from Rodrigue's formula

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} [(x^2 - 1)^n]$$

2. Obtain the Legendre polynomial  $P_4(x)$  directly from Legendre's equation of order 4 by assuming a polynomial of degree 4, i.e.

$$y = ax^4 + bx^3 + cx^2 + dx + e$$

3. Obtain the Legendre polynomial  $P_6(x)$  by application of the recurrence formula

$$nP_n(x) = (2n - 1)xP_{n-1}(x) - (n - 1)P_{n-2}(x) \text{ assuming that } P_4(x)$$

and  $P_5(x)$  are known.

4. Obtain the Legendre polynomial  $P_2(x)$  from Laplace's integral formula

$$P_n(x) = \frac{1}{\pi} \int_0^\pi (x + \sqrt{x^2 - 1} \cos t)^n dt$$

5. Find the first three coefficients in the expansion of the function

$$\sqrt[3]{1 - x} \quad -1 \leq x \leq 1$$



$$f(x) = \begin{cases} x & -1 \leq x < 0 \\ 0 & 0 \leq x < 1 \\ x & 1 \leq x \leq 1 \end{cases}$$

in a series of Legendre polynomials  $P_n(x)$  over the interval  $(-1, 1)$ .

6. Find the first three coefficients in the expansion of the function

$$f(\theta) = \begin{cases} \cos \theta & 0 \leq \theta \leq \pi/2 \\ 0 & \pi/2 \leq \theta \leq \pi \end{cases}$$

in a series of the form

$$f(\theta) = \sum_{n=0}^{\infty} A_n P_n(\cos \theta) \quad 0 \leq \theta \leq \pi$$

7. Obtain the associated Legendre functions  $P_2^1(x)$ ,  $P_3^2(x)$  and  $P_2^3(x)$ .
8. Verify that the associated Legendre function  $P_3^2(x)$  is a solution of Legendre's associated equation for  $m = 2$ ,  $n = 3$ .
9. Verify the result

$$\int_{-1}^1 P_n^m(x) P_k^m(x) dx = 0 \quad n \neq k$$

for the associated Legendre functions  $P_2^1(x)$  and  $P_3^1(x)$ .

10. Verify the result

$$\int_{-1}^1 [P_n^m(x)]^2 dx = \frac{2}{2n+1} \frac{(n+m)!}{(n-m)!}$$

for the associated Legendre function  $P_1^1(x)$ .

11. Obtain the Legendre functions of the second kind  $Q_0(x)$  and  $Q_1(x)$  by means of

$$Q_n(x) = P_n(x) \int \frac{dx}{[P_n(x)]^2(1-x^2)}$$

12. Obtain the function  $Q_3(x)$  by means of the appropriate recurrence formula assuming that  $Q_0(x)$  and  $Q_1(x)$  are known.

